


201-146
FEED TO ANIMALS



VAPONA® INSECTICIDE
EMULSIBLE CONCENTRATE

(Contains 10.0 lbs. VAPONA Insecticide per gal.)

ACTIVE INGREDIENTS	BY WEIGHT
2,2-dichlorovinyl dimethyl phosphate.....	81.2%*
Related Compounds.....	6.3%
INERT INGREDIENTS.....	<u>12.5%</u>
TOTAL.....100.0%	

*Equivalent to 87.5% w VAPONA Insecticide.

USDA REG. NO. 201-146

GENERAL DIRECTIONS

VAPONA® Insecticide Emulsible Concentrate (10 pounds per gallon) is a specially prepared concentrate for use in conventional application equipment.

USE AND DIRECTIONS FOR APPLICATION
(INSERT ATTACHED USES HERE)

DANGER

KEEP OUT OF REACH OF CHILDREN, DOMESTIC ANIMALS AND PETS.

POISON



PRECAUTIONS IN USING

VAPONA Insecticide concentrate and the diluted solution are poisonous. Therefore precautions must be observed when handling them. (See antidote and warning below).

Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted solution. It is recommended that only the amount of diluted solution be prepared from the concentrate that will be used immediately.

Do not contaminate lakes and streams.

Do not contaminate feed, water or foodstuffs.

SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING
STATEMENTS ON SIDE PANEL.

NET CONTENTS _____

SIDE PANEL

WARNING

VAPONA Insecticide concentrate is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid prolonged exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare dilutions. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for DDVP (VAPONA Insecticide). Keep all unprotected persons out of operational area.

In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

ANTIDOTE AND FIRST AID TREATMENT

Atropine is the emergency antidote for VAPONA Insecticide poisoning. Consult your Physician about obtaining an adequate supply of 1/100-grain atropine for emergency use.

Call a physician immediately in all cases of suspected poisoning.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water).

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately.

NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below).

KEEP PATIENT PRONE AND QUIET.

START ARTIFICIAL RESPIRATION IMMEDIATELY IF PATIENT IS NOT BREATHING.

TRANSPORT THE PATIENT IMMEDIATELY TO THE NEAREST PHYSICIAN.

NOTE TO PHYSICIAN

diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

Morphine is contraindicated.

Clear chest by postural drainage.

Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

Allow no further exposure until blood tests indicate that cholinesterase levels have returned to normal.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

SHELL CHEMICAL COMPANY
A DIVISION OF SHELL OIL COMPANY
AGRICULTURAL CHEMICALS DIVISION
New York, New York 10020

CROP OR USE	INSECT	DOSAGE ACTUAL VAPONA [®] INSECTICIDE	DIRECTIONS FOR APPLICATION	INTERVAL, FINAL TREATMENT TO HARVEST. DAYS
Mush- rooms	Sciarid Flies, Phorid Flies	32 grams per 16,000 cu.ft. EC, Aerosol	Apply in an Aerosol or hot pipe method (0.9 fl.oz. of 10 lb/gal solvent-free EC per 16,000 cu.ft. placed in a hot plate or steam pipe to vaporize the VAPONA) every 4 days as a preventive treat- ment or when insect infesta- tions are present. Repeat as often as necessary to main- tain control. During applica- tion, use a mask or respirator of a type passed by the U. S. Department of Agriculture for DDVP (VAPONA [®] Insecticide) protection. Mushroom houses should be kept closed for several hours after treatment and workers should not re- occupy the premises until it has been ventilated.	1 (one)