



SOLID RED

AZODRIN[®] 5

INSECTICIDE

DO NOT STORE AT TEMPERATURES BELOW 70°F OR ABOVE 80°F FOR PROLONGED PERIODS.



DANGER! POISON
KEEP OUT OF THE
REACH OF CHILDREN



Do not store AZODRIN 5
Water Miscible Insecticide
at temperatures below 70 F
or above 80 F
for prolonged periods.



Do not store AZODRIN 5
Water Miscible Insecticide
at temperatures below 70°F
or above 80°F
for prolonged periods.

for at least ten minutes and get medical attention. In case of contact, re-
move clothing and wash skin with soap and water. Wash clothing before
reuse. Swallowing of AZODRIN 5 may cause dizziness, nausea, vomiting, and
stomach pain. Swallowing of large amounts may cause unconsciousness. These
symptoms occur and are relieved by washing with a dilute solution of alkali
then with 1% acid solution and then with water.

NOTE: AZODRIN 5 should also be used on ornamentals when handling ornamental
containers.

PREVENTION OF CONTAMINATION AND OTHER USES: Do not reuse for any
purpose. Decontaminate containers by washing thoroughly with a dilute
solution of alkali (less than 5%) and detergent and rinsing with water.
Empty containers by crushing and burying fragments in an isolated area.
Dispose of rinsings in a way so not to constitute a hazard, or contaminate
water supplies.

ANTIDOTE AND FIRST AID TREATMENT

**AZODRIN 5 IS THE EMERGENCY ANTIDOTE FOR AZODRIN INSECTICIDE
POISONING.** AZODRIN 5 is of no antidote and may be used in conjunction with
atropine. CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.

SYMPTOMS: If the material has been swallowed, induce vomiting immedi-
ately. This may be done by introducing a finger into the throat or by giving
warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE
ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

SYMPTOMS: If the material has been spilled on the skin, immediately re-
move patient from the vicinity of the insecticide, remove all contaminated
clothing, and wash skin with soap and running water. If the material gets
into the eyes, wash immediately with running water for at least ten min-
utes. IF SERIOUS SYMPTOMS APPEAR: (See WARNING SYMPTOMS below) Keep patient
prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest
hospital.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness
in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating,
nausea, vomiting, diarrhea, and abdominal cramps.

PREVENTION: Atropine is the specific therapeutic antagonist of cholinergic
stimulation. If there are signs of para-
sympathetic stimulation, atropine sulfate should be injected at two-minute
intervals, in doses of 1 to 2 milligrams, until complete atropinization has
occurred.

Pralidoxime chloride (2-PAM chloride) may also be used as an effective
antidote in addition to and while maintaining full atropinization. In adults,
an initial dose of 1 gram of 2-PAM should be injected, preferably as an in-
fusion in 250 cc of saline over a 15- to 30-minute period. If this is not
practical, 2-PAM may be administered slowly over not less than two min-
utes. After about an hour, a second dose of 1 gram of 2-PAM will be in-
dicated if muscle weakness has not been relieved. For infants and children,
the dose of 2-PAM is 0.25 gram.

Morphine is an improper treatment.

Clear chest by postural drainage. Oxygen administration may be necessary.
Observe patient continuously for 48 hours. Repeated exposure to cholin-
esterase inhibitors may, without warning, cause prolonged susceptibility
to very small doses of any cholinesterase inhibitor. Allow no further ex-
posure until cholinesterase regeneration has been obtained as determined
by blood test.

NOTICE OF WARRANTY

SMALL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY,
FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, con-
cerning this product or its use which extend beyond the use of the product
under normal conditions in accord with the statements made on this label.

GROUP	PLANT OR ANIMAL	RATE Gallons/Acre	USE INSTRUCTIONS	USE RESTRICTIONS
Cotton	Plant Bugs & Eggs	1/2 to 1 1/2	For early season control use 1/2 gal. per acre. For high populations, use 1 to 1 1/2 gal. per acre.	Do not apply within 21 days of harvest. Do not graze livestock on treated fields or feed treated trash.
	Bollworms, Pink Bollworms, Loopers, Bolls, Boll Weevils	1/2 to 1 1/2	To control light infestations use 1/2 gal. per acre. For higher populations use 1 to 1 1/2 gal. per acre. One gallon will treat 5 acres at the 1 1/2 gal./acre dosage. Apply to give uniform coverage. Repeat as necessary to maintain control but not more frequently than every 5-7 days.	
	Thrips	3-5 ozs		
	Beet Armyworm	1/2 to 1 1/2		
Peanuts	White Flies	1/2 to 1 1/2		NOTE: California-San Joaquin Valley Only: Do not apply later than 2 weeks after initiation of bloom and definitely not after July 15 in order to avoid injury to wildlife.
	Lesser Cornstalk Borer	1 1/2	Apply the spray in a 10 to 12 inch band which covers only the soil surface and lowest stems and foliage.	
	Worms (Southwestern United States)	1 1/2 to 1 1/2		
	Worms (Southeastern United States)	1/2 to 1 1/2	Use the lower dosage for light infestations and the higher dosage for heavy infestations.	
Sugarcane	Granulate Cutworms	1 1/2		Repeat applications as necessary to maintain control, applying no more than 2 applications per season. Do not feed treated vines or hay to livestock. Do not graze livestock on treated fields.
	Sugarcane Borers	1-1 1/2	For aerial applications use 19.2 ozs. in minimum of 2 gals. of water per acre. To insure uniform coverage aircraft applications should be made as close as possible to canopy level. Repeat applications as necessary to maintain control.	
	FLORIDA ONLY: Aphids, Mites			
Tobacco	Tobacco Hornworm	1/2	Apply with sufficient water to give uniform coverage. Where specified, use lower dosage for light infestations and the higher dosage for heavy infestations. Repeat as necessary to maintain control.	Do not use on shade grown tobacco. Do not apply within 5 days of harvest.
	Green Peach Aphid	1/2		
	Tobacco Budworm	1/2 to 1 1/2		
Potatoes (White, Irish)	Colorado Potato Beetle, Leafhoppers	1/2 to 1 1/2	Apply in not more than 100 gallons of water to give uniform coverage. Where specified, use the lower dosage for light infestations and the higher dosage for heavy infestations. Repeat application as necessary to maintain control.	Do not apply within 7 days of harvest.
	Fire Beetle	1/2		
	Aphids	1/2 to 1 1/2		
	Worms (Southwestern United States)	1/2 to 1 1/2		
	Potato Tuberworm	1 1/2		
Tomato (FLORIDA ONLY)	Aphids	1/2	Apply with ground equipment only. Apply with sufficient water to give uniform coverage. Applications should begin at first appearance of insect populations. Repeat as necessary to maintain control.	Do not apply within 21 days of harvest.
	Leafminers, Cabbage Loopers, Tomato Fruitworm	1 1/2		

CODE 2000 JG