ACCEPTED

FEB 1 0 1975

Under the Feder Insecticide. Fungicial, and Rodentistide Act. as amended, for the pesticiae registered under EPA Reg. No.

> DIRECTIONS: Place Flea Collar around dog's neck, adjust for proper fit and snap in place. The collar must be worn loosely and in addition to the regular collar. If collar is worn too tightly, it may produce neck irritation. Cut off and dispose the excess length. Flea kill begins at once and will continue for 3 months while the collar is worn. Ticks are harder to kill but those on the neck area will soon be killed and easily removed. Collar will continue to kill ticks for eight weeks. For continuous protection, replace collar every 3 months.

> flea Collar can be worn with regular collar. Remove collar when washing dogreplace when dry.

> CAUTION: Do not open protective pouch until ready to use. Do not allow children to play with this collar. Collar is intended for use only as an insecticide generator (Shell's U.S. Patents 2,956,073, 3,116,201 and 3,318,769) and is not to be taken internally by man or animals. Do not use on cats, whippets and greyhounds. Some dogs may be sensitive to this collar. Remove collar at first sign of insitation or adverse reaction. Do not use on sick or convalescing dogs. Do not use any other pesticide on dog while collar is worn.

> > Manufactured by SHELL CHEMICAL COMPANY A Division of Shell Oil Company ANIMAL HEALTH SAN RAMON, CA. 94583

EPA REG. NO.201-230 AA **EPA EST 201-NJ-1**





BLACK

FOR LONG-LASTING FLEA CONTROL AIDS IN TICK CONTROL — ESPECIALLY IN THE NECK AREA **Veterinary Product**

CAUTION:

Do not allow children to play with this collar — see cautions on back panel.

ACTIVE INGREDIENTS BY WEIGHT 2,2-Dichloroviny! dimethy! phosphate *Equivalent to 10% w TECHNICAL VAPONA® Insecticide

CONTENTS: 1 FLEA COLLAR

NET WEIGHT: 0.8 OZ.

ACF 217D