



AKTON[®] INSECTICIDE

TECHNICAL (SD 9098)
ORGANIC PHOSPHATE COMPOUND, LIQUID N.O.S.

FOR MANUFACTURING PURPOSES ONLY

ACTIVE INGREDIENTS	BY WEIGHT
O-(2-chloro 1-(2,5-dichlorophenyl) vinyl)	80%
O,O-diethyl phosphorothioate (MP 27°C)	
Related compounds	5%
INERT INGREDIENTS	15%
TOTAL	100%

WARNING  **POISON**

KEEP OUT OF THE REACH OF CHILDREN

HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing and clean rubber gloves. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. If swallowed, induce vomiting immediately and get medical attention. Keep away from heat and open flames. Do not contaminate fish-bearing waters by cleansing of equipment or disposal of wastes.

DECONTAMINATE AND DESTROY USED CONTAINERS: Do not re-use for any purpose. Decontaminate containers by washing thoroughly with a strongly alkaline detergent solution and rinsing with water. Dispose of rinsing in a way as not to constitute a hazard or contaminate water supplies.

Destroy containers by crushing or puncturing; then bury, or burn; if burned stay out of smoke.

Note: Caution! Avoid skin and eye contamination when handling strong caustic solutions.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR VAPONA INSECTICIDE POISONING. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR AKTON INSECTICIDE POISONING. Consult your physician about obtaining an adequate supply of 1/100-grain atropine tablets for emergency use.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately.

NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below). Keep patient prone and quiet.

Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.


NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against para-sympathetic nervous stimulation. If there are signs of para-sympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

SHELL CHEMICAL COMPANY

A DIVISION OF SHELL OIL COMPANY, NEW YORK, N. Y.



NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.