diarrhea and abdominal cramps.

Λ\_

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, Indoses of 1 to 2 milligrams until complete atropinization has occurred.

Morphine is contraindicated.

Clear chest by postural drainage.

Observe patient continuously for 48 hours.

Repeated exposure to cholinestrase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinestrase inhibitor.

Allow no further exposure until blood tests indicate that cholinesterase levels have returned to normal.

# NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

SHELL CHEMICAL COMPANY
A DIVISION OF SHELL OIL COMPANY
AGRICULTURAL CHEMICALS DIVISION
New York, New York 10020

CROP DOSAGE ACTUAL OR VAPONA® USE INSECT INSECTICIDE	DIRECTIONS FOR APPLICATION	INTERVAL, FINAL TREATMENT TO HARVEST, DAYS
--	----------------------------	--

Mush- Sciarid 32 grams per rooms Files, 16,000 cu.ft. Phorid EC, Aerosol Flies

Apply in an Aerosol or hot pipe method (0.9 fl.oz. of 10 lb/gal solvent-free EC per 16,000 cu.ft. placed in a hot plate or steam pipe to vaporize the VAPONA) every 4 days as a preventive treatment or when insect infestations are present. Repeat as often as necessary to maintain control. During application, use a mask or respirator of a type passed by the U.S. Department of Agriculture for DDVP (VAPONA® Insecticide) protection. Mushroom houses should be kept closed for several hours after treatment and workers should not reoccupy the premises until 1t has been ventilated.

1 (one)

SHELL CHFITCAL COMPANY
A Division of Shell Oil Company
Agricultural Chemicals Division

New York, New York 10020

1(0f2)



ACCEPTED.

APR 2 1969

# VAPONA(R) INSECTICIDE 50% CONCENTRATE For Manufacturing Purposes Only

ACTIVE INGREDIBITS

BY WEIGHT

2,2-Dichlorovinyl dimethyl phosphate Related Compounds Petroleum hydrocarbons 46.5%\* 3.5%\* 50.0% 100.0%

TOTAL

\* Equivalent to 50%w VAPONA(R) Insecticide

USDA REG. NO.

Net Contents



#### PRECAUTIONS IN USING

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. PCISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES.

Do not get on skin or in eyes. Do not breathe fumes. Avoid prolonged exposure to fumes. Wear clean rubber gloves, protective clothing and goggles when handling the concentrate to prepare dilutions. During use indoors wear a full face mask or a type passed by the U. S. Department of Agriculture for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational area. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted solution. Do not conteminate feed, water or foodstuffs.

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

DECONTAMINATE AND DESTROY USED CONTAINEDS: Do not re-use for any purpose. Decontaminate containers by washing thoroughly with a strongly alkaline detergent solution and rinsing with water. Dispose of rinsings in a way as not to constitute a hazard or contaminate water supplies.

Destroy containers by crushing or puncturing; then bury, or burn: if burned stay out of smoke.

Note: Caution! Avoid skin and eye contamination when handling strong caustic solutions.

2(0/3)

### ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR VAPONA INSECTICIDE POISONING. Consult your physician about obtaining an adequate supply of 1/100-grain atrovine tablets for emergency use. CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). Repeat warm salt water until vomit fluid is clear. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Have patient lie down and keep quiet.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

#### NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

# NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

SOLID RED

PRECAUTIONS IN USING

ATTACHMENT OF A DIRECTOR A DIRECTOR A DIRECTOR AND ADDRESS OF A DIRECTOR ADDRESS OF A DIRECTOR AND ADDRESS OF A DIRECTOR ADDRESS OF ADDRESS OF A DIRECTOR ADDRESS OF ADDRESS OF A DIRECTOR ADDRESS OF ADDRESS OF ADDRESS OF A DIRECTOR ADDRESS OF ADDRES

SAN RAMON, CALIFORNIA 94583

# VAPONA

**EMULSIFIABLE-**

FLY and MOSQUITO CONTROL:
ON DAIRY and BEEF CATTLE

IN and AROUND ANIMAL BUILDINGS

ACTIVE INGREDIENTS

2,2 Dichlorovinyl dimethyl phosphate 21.8%\*

Related Compounds

1.6%\*

Petroleum Hydrocarbons
INERT · INGREDIENTS

1.6%\* 71.6% 5.0%

100.0%

TOTAL

DANGER!

KEEP OUT OF THE REACH OF CHILDREN

SEE ANTIDOTE AND OTHER PRECAUTIONS ON SIDE PANELS

EPA REG. NO. 201-158 ZC

MUST BE DILUTED

TAKE TIME

OBSERVE LABEL


DIRECTIONS

\*Equivarent to 23.4% w VAPONA\* Insecticide

NET CONTENTS: 1 GALLON



-SOL'D RE CENTER



#### ANTIDOTE AND FIRST AID TREATMENT-

ATROPINE is the emergency antidote for VAPONA Insecticide poisoning. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERMAL: If the material has beer, swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of wat ... NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

**EXTERMAL:** If the material has sen spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least 10 minutes.

#### IF WARNING SYMPTOMS APPEAR:

Keep patient prone and quiet.

Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital. (See WARNING SYMPTOMS below.)

#### NOTE TO PHYSICIAN-

WARRING SYMPTOMS: Symptoms include weakness, headache tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

#### Morphine is contraindicated.

Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization. In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion in 250 cc of saline over a 15 to 30 minute period. If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5 percent solution in water over not less than two minutes. After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved. For infants and children the dose of 2-PAM is 0.25 grams.

# Crear chest by postural drainage.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

#### -PRECAUTIONS IN USING

YAPONA Insecticide Emulsifiable and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings statement.)

Store VAPORA Insecticide Emulsifiable in a safe place. Children or animals must not be able to reach or handle VAPORA Insecticide Emulsifiable or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from VAPORA Insecticide Emulsifiable. Not for use or storage in or around the home.

Mix YAPONA Insecticide Emulsifiable into water outdoors or in a well ventilated room, Wash off the outside of the bettle of YAPONA Insecticide Emulsifiable in running water before storing away.

This product is texic to fish, birds and wildlife. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weether conditions favor drift from areas treated. Apply this product only as specified on this label. Do not contaminate feed, water or foodstuffs, milk or milking utensils.

#### -DANGER

VAPONA Insecticioe Emulsifiable is peisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid exposure to fumes. Wear clean rubber gloves, waterproof protective clothing and goggles when handling the VAPONA Insecticide Emulsifiable to prepare emulsions. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DOVP (VAPONA Insecticide) protection. Keep all unpretected persons out of eperational area.

In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

#### DECONTAMONATE AND DESTROY USED CONTAMERS.

Decontaminate containers by washing thoroughly with a strong solution of alkali and detergent and rinsing with water.

Destroy containers by crushing or puncturing; then bury or burn.

Dispose of rinsings in a way as not to constitute a hazard, or contaminate water

#### -NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES HO WARRANTY OF MERCHANTABILITY, FITHESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

# VAPONA°

EMULSIFIABLE-

FLY and MOSQUITO CONTROL: ON DAIRY and BEEF CATTLE

IN and AROUND ANIMAL BUILDINGS

DANGER! POISO

MUST BE DILUTED

ANNIAL HEALTH

VAPONA on use, knapsaci percent the safe amount wapona when us

DIRECT

DIRECT Dairy Beef

FOGGING Dairy Pourtr Horse Suring Other Build

FOSGING Cattle Stocky Holdid Corra Other Anie MHELL CHATTERE C. MALY

A Divinion, Constituted transArricalta - Constitute Division.

New York, New York 100 c

VAPONA(F) INCHMENTATION CONCENTRATES
For Manufacturing Purposes Only

1.21

ACTIVE INCR. DID NO.

UMA MO. H.

BY WEIGHT

2,2-Dichloroving 1 directly 1 phosebute Related Conscions

50.0%

Petroleum hydrocari ona

TATAL

\* Equivalent to M to VANTAA(I) Insecticities

Net Contents .....

DANGIN

ióin a

THE POWER OF THE PARTY OF THE PARTY

THE RESIDENCE OF THE PROPERTY OF THE PROPERTY

In not pet in Alin or in even, to bet the the face . As it brokens, in a consideration of the the consequent of a consideration of the constant of a constant of

or kan an kanan segarah kembendagi mengantan geberakan debah beranda sebenah beranda debenah di beranda di beb Geberah di beginan sebenah di beranda di beranda di beranda di beberakan di beranda di beranda di beranda di b Geberah genah genah di beranda

engan sa mengan kenalagan di sebagai kenalagan sebagai di kenalagan sebagai kenalagan sebagai kenalagai kenala Sebagai kenalagai sebagai seba

i de la compartición de la compa

en de la companya de la co

# ANTIBOTE AND FIRST ARE TO EASTITHT

ATROPINE IS THE EXHIBITION ANTHOUGH WAS A TOO FIGURE POLICE OF COnsult your physician about obtaining an AC pate analy of 1/100-grain atropine tellets for emergency . A A PHYSICIAN IN ALL WARD OF SUPPRETED POISONING.

INTrans. ... material has been suallowed, in bace veniting immediately. This may be do not by introducing a finger into the threat or by giving warm salt water (I tablespoon of salt to a glass of water). Repeat warm salt water until venit fluid is clear. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Have patient lie down and keep quiet.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100-grain atropine tablets immediately. NEVER ADMINISTUR ATROPINE UNLESS WARNING SYMPTOMS APPEAR. (See WARNING SYMPTOMS below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

### NOTE TO PHYSICIAN

WARRING SYMPTOMS: Symptoms include weakness, headache, tichtness in chect, llurred vision, non-reactive pin-roint pubils, salivation sweating, nauges, veniting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therareutic antatonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 15 minute intervels, in doses of 1 to 2 milligrams, until complete atropinisation has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 43 hours. Repeated exposure to cholinesterase inhilitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhilitors. Allow no further exposure until cholinecterase regeneration has been attained as determined by blood test.

# TRAIN OF WHATER

Child Childed data in the in the second of the content of the content of the second of the uses which extend topical the use of the crossic order rough exceptions to second with the stellownite made on this label.