

DIRECTIONS FOR USE (continued)

ORNAMENTALS: continued-

Azalea Scale, Bagworm, Birch Leaf Miner, Boxwood Leaf Miner, Fletcher Scale, Florida Red Scale, Juniper Scale, Magnolia Scale, Oak Kermes, Pine Leaf Scale, Tent Caterpillar - Use 2 pints per 100 gals., dilute spray (2 teaspoonfuls per gal.). For Fletcher, Juniper and Oak Kermes Scales, apply when scale crawlers have settled on foliage.

CAUTION: Avoid use on certain ferns including Boston, Maidenhair and Pteris, as well as on some species of Crassula.

GRASSHOPPERS

WASTELANDS, ROADSIDES, SOIL-BANK LANDS (Not Grazed): Use 1½ to 3 pts. per acre in water for good coverage or in 1 gal. diesel fuel oil per acre by means of an airplane or a turbine-blower type sprayer.

FLY CONTROL

FLY CONTROL IN BUILDINGS: For use in and around buildings which house domestic animals, around the outside of homes, around and in non-edible product areas of meat processing plants.

STRAIGHT MALATHION SPRAYS		MALATHION BAIT SPRAYS			
Amount of Spray	Amount of Malathion S-E	Amount of Bait Spray	Amount of Malathion S-E	Sugar (or) Unsulphurized Molasses or Corn Syrup	Unsulphurized Molasses or Corn Syrup
2½ gals.	1 cup	2½ gals.	1 cup	1 cup	1 cup
12 gals.	1 qt.	12 gals.	1 qt.	2½ lbs.	1 qt.
100 gals.	2 gals.	100 gals.	2 gals.	20 lbs.	2 gals.

Apply the spray at the rate of 1 gal. per 1,000 sq. ft. on painted surfaces and 2 gals. per 1,000 sq. ft. on unpainted surfaces where flies alight or congregate. Sugar, unsulphurized molasses, or corn syrup have been found in most cases to prolong the insecticidal activity of Malathion against flies. Do not apply Malathion within 14 days to freshly whitewashed surfaces.

CAUTION: Avoid contamination of milk, milk equipment and water. Avoid contamination of feed and food products, also drinking fountains and feed troughs. Remove calves under one month of age and lactating dairy animals from buildings before treating. Do not leave within reach of children.


STORED GRAINS: For the protection of stored wheat, oats, corn, rye, barley grains, grain sorghum, and field or garden seeds, against confused flour beetle, rice weevil, granary weevil, saw-toothed grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer, Indian meal moth, and for control of cereal leaf beetles, apply CYTHION emulsifiable liquid as follows:

RESIDUAL SPRAY-BEFORE STORING GRAINS: For a residual wall, floor, and machinery spray in grain elevators, in treating truck beds, box cars, and ships-holds before loading grain, apply 1 gallon of CYTHION emulsifiable liquid per 25 gals. of water making thorough application. Before applying spray, clean elevators, box cars, etc. thoroughly. Remove and burn all sweepings and debris. Wear a mask or respirator of type recommended by the U.S. Bureau of Mines to give adequate protection against cythion. Wear protective clothing to minimize skin exposure.

GRAINS GOING INTO STORAGE: Apply 1 pint of CYTHION emulsifiable liquid in 2-5 gallons of water per 1000 bushels. Apply as the grain is being loaded or turned into final storage.

AFTER GRAINS ARE STORED: To protect stored grains from attack by Indian meal moth, apply CYTHION emulsifiable liquid to the surface of clean or uninfested grain at the rate of ½ pint in 1-2 gals. of water per 1000 sq. ft. of grain surface area. Apply the spray evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

ACCEPTED
201-271
5-26-71
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
ECONOMIC POISON REGISTERED
UNDER NO.



ADD-1850 2-71 BEND TAB TO PEEL LABEL

NET CONTENTS: 29 GAL.
NET WEIGHT: 299 LBS.

Accepted With Comments

FEDERAL REGISTER
DEPARTMENT OF AGRICULTURE
Pesticide Registration Division

ADD-1850 2-71 BEND TAB TO PEEL LABEL



NET CONTENTS: 29 GAL.

NET WEIGHT: 299 LBS.