

REFER TO SUPPLEMENTAL LABELING FOR DIRECTIONS ON THE USE OF BLADEX HERBICIDE IN TANK MIX COMBINATIONS

WEEDS CONTROLLED BY BLADEX 4 WDS HERBICIDE ALONE AND IN COMBINATION WITH HERBICIDE TANK MIXES SHOWN IN THE SUPPLEMENTAL LABELING

GRASSES

Barrygrass Barnyardgrass Digitaria Johnsongrass Panicum	Redtop Ryegrass St. Augustine Ternstroemia Timothy Wheatgrass	Green foxtail Hempdog Mile-a-minute Spikegrass	Green foxtail Hempdog Mile-a-minute Spikegrass	Stinkgrass Watergrass Yellow foxtail
---	--	---	---	--

BROADLEAVES

Blackberry Cottonwood Dandelion Hemp Marestail Ragweed Rasp Sickle-leaved Soybean Tack Tackgrass Tobacco Waterhemp Witchweed Yellow flowered	Blackberry Cottonwood Dandelion Hemp Marestail Ragweed Rasp Sickle-leaved Soybean Tack Tackgrass Tobacco Waterhemp Witchweed Yellow flowered	Marestail Ragweed Rasp Sickle-leaved Soybean Tack Tackgrass Tobacco Waterhemp Witchweed Yellow flowered	Plantain Puncture Ragweed Rasp Sickle-leaved Soybean Tack Tackgrass Tobacco Waterhemp Witchweed Yellow flowered Yellowweed-Pennsylvanian	Yarrow Sunflower Tarnweed Tobacco Waterhemp Witchweed Yellow flowered Yellowweed-Pennsylvanian
--	--	---	--	---

GENERAL INFORMATION

Bladex 4 WDS Herbicide is a selective herbicide which controls many annual and perennial broad-leaved weeds in corn, sorghum, soybeans, cotton, and grain sorghum. It is also effective on some grasses and sedges. Bladex 4 WDS Herbicide is a water dispersible suspension. It is applied by spray or broadcast application.

Bladex 4 WDS Herbicide is a selective herbicide which controls many annual and perennial broad-leaved weeds in corn, sorghum, soybeans, cotton, and grain sorghum. It is also effective on some grasses and sedges. Bladex 4 WDS Herbicide is a water dispersible suspension. It is applied by spray or broadcast application.

Bladex 4 WDS Herbicide is a selective herbicide which controls many annual and perennial broad-leaved weeds in corn, sorghum, soybeans, cotton, and grain sorghum. It is also effective on some grasses and sedges. Bladex 4 WDS Herbicide is a water dispersible suspension. It is applied by spray or broadcast application.

Bladex 4 WDS Herbicide is a selective herbicide which controls many annual and perennial broad-leaved weeds in corn, sorghum, soybeans, cotton, and grain sorghum. It is also effective on some grasses and sedges. Bladex 4 WDS Herbicide is a water dispersible suspension. It is applied by spray or broadcast application.

APPLICATION DIRECTIONS

GENERAL MIXING AND SPRAYING DIRECTIONS

Bladex 4 WDS Herbicide should be mixed with clean water. Do not use Bladex 4 WDS Herbicide on sands or heavy sands and do not use on soil with low water-holding capacity. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

Bladex 4 WDS Herbicide should be mixed with clean water. Do not use Bladex 4 WDS Herbicide on sands or heavy sands and do not use on soil with low water-holding capacity. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

Bladex 4 WDS Herbicide should be mixed with clean water. Do not use Bladex 4 WDS Herbicide on sands or heavy sands and do not use on soil with low water-holding capacity. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

Bladex 4 WDS Herbicide should be mixed with clean water. Do not use Bladex 4 WDS Herbicide on sands or heavy sands and do not use on soil with low water-holding capacity. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

PREFEMERGENCE

Bladex 4 WDS Herbicide should be applied to the soil before the weed seedlings emerge. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

Bladex 4 WDS Herbicide should be applied to the soil before the weed seedlings emerge. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

Bladex 4 WDS Herbicide should be applied to the soil before the weed seedlings emerge. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

PREFEMERGENCE BROADCAST APPLICATION RATES FOR BLADEX 4 WDS

For broadcast application use proportionately less.

TYPE OF APPLICATION	PERCENT ORGANIC MATTER IN SOIL*					
	Less than 1%	1%	2%	3%	4%	Over 4%
	Quarts of BLADEX 4 WDS Per Acre	Quarts of BLADEX 4 WDS Per Acre	Quarts of BLADEX 4 WDS Per Acre	Quarts of BLADEX 4 WDS Per Acre	Quarts of BLADEX 4 WDS Per Acre	Quarts of BLADEX 4 WDS Per Acre
Preemergence Broadcast	1.0	1.5	2.0	2.4	2.8	3.2
Preemergence Broadcast	1.0	1.5	2.0	2.4	2.8	3.2
Preemergence Broadcast	1.0	1.5	2.0	2.4	2.8	3.2
Preemergence Broadcast	1.0	1.5	2.0	2.4	2.8	3.2
Preemergence Broadcast	1.0	1.5	2.0	2.4	2.8	3.2

PRECAUTIONS IN USING

Bladex 4 WDS Herbicide should be used in accordance with the directions on the label. Do not use Bladex 4 WDS Herbicide on soils which are highly acidic or highly alkaline.

↑ **THIS SIDE UP** ↑



Bladex 4
WATER DISPERSIBLE SUSPENSION

Bladex 4
WATER DISPERSIBLE SUSPENSION

CONTENTS OF THIS LABEL
This label contains the directions for the use of Bladex 4 WDS Herbicide. It is a selective herbicide which controls many annual and perennial broad-leaved weeds in corn, sorghum, soybeans, cotton, and grain sorghum. It is also effective on some grasses and sedges. Bladex 4 WDS Herbicide is a water dispersible suspension. It is applied by spray or broadcast application.

SHELL CHEMICAL COMPANY

ACD-129F 12-75

SIDE UP

TO OPEN

Insert thumb in notch.
Pull back toward handle.
Separate from unit.




↑ THIS SIDE UP ↑



Bladex[®] 4
WATER DISPERSIBLE SUSPENSION
Herbicide



Bladex[®] 4
WATER DISPERSIBLE SUSPENSION
Herbicide



Bladex[®] 4

WATER DISPERSIBLE SUSPENSION



Bladex[®] 4

WATER DISPERSIBLE SUSPENSION
Herbicide

(lb. Bladex per gallon)

	BY WEIGHT
.....	43%
.....	57%
TOTAL	100%

SEE INSTRUCTIONS FOR USE AND WARRANTY BEFORE OPENING
THIS PRODUCT IS NOT ACCEPTABLE
IF THE CONTENTS ARE NOT AS SHOWN

WARNING!
KEEP OUT OF REACH OF CHILDREN.
See side panels for additional precautions.

NET CONTENTS: 5 GALLONS

SEE INSTRUCTIONS FOR USE AND WARRANTY BEFORE OPENING THIS PRODUCT IS NOT ACCEPTABLE IF THE CONTENTS ARE NOT AS SHOWN

**CODE
4360MB**

1801-00-51

GREY AREA IS SOLID GREEN
LABEL SHOWN IS SIZE
ALL COPY LINE IS OTHERWISE INDICATED IS SOLID BLACK

SUPPLEMENTAL LABEL FOR USE OF BLADEX® 4 WDS HERBICIDE IN TANK MIXES WITH ATRAZINE, ALACHLOR

(Refer to label on container for ingredients, precautions, notice of warranty and use directions for BLADEX® alone and in tank mixes)

GENERAL INFORMATION

BLADEX 4 WDS Herbicide is a selective preemergence or postemergence herbicide for the control of annual grasses and broadleaf weeds in field corn. BLADEX 4 WDS Herbicide may also be used preemergence in certain tank mix combinations with other herbicides as described below.

When BLADEX 4 WDS Herbicide is used alone or in combination with Sutan, Alachlor, Lasso, or Paraquat as directed in this label, the user can safely plant corn in other crops. When BLADEX 4 WDS Herbicide is used in combination with atrazine, however, only corn or sorghum may be planted the following year.

As a preemergence herbicide, BLADEX 4 WDS Herbicide is active mainly through the roots and therefore its effectiveness is dependent on adequate rainfall to move the herbicide into the root zone. A rotary hoeing or shallow cultivation is recommended if a rainfall or sprinkler irrigation has not occurred within about five days after application of BLADEX 4 WDS Herbicide alone, as a tank mix or in a sequential application. Moisture should be sufficient to wet the top 2 to 3 inches of soil to make the soil too wet to cultivate; for most soils one half inch of moisture is sufficient. Under conditions which dry weeds or plants, such as a few perches, lack of soil surface moisture, or when germination extends over a long period, the effectiveness of the herbicide may be impaired. Supplementary cultural practices may be of benefit under these circumstances.

BLADEX 4 WDS Herbicide is not effective when used preemergence on peat or muck soils. Do not use BLADEX 4 WDS Herbicide on sands or heavy sands (soils consisting of more than 70% sand) containing less than 1% organic matter. BLADEX 4 WDS Herbicide should be applied only once per crop season.

Observe all cautions and limitations on labeling of all products used in mixtures. The effectiveness of any combination is generally dependent on weed spectrum, soil texture, and soil organic matter content. Suggested rates for all tank mix combinations of BLADEX 4 WDS Herbicide with other herbicides can be found in rate tables contained in this label and in supplemental literature.

APPLICATION DIRECTIONS

GENERAL MIXING AND SPRAYING DIRECTIONS

Be sure the sprayer tank, boom and nozzles have been thoroughly flushed with clean water if another pesticide has been used previously. Sufficient jet or mechanical agitation should be provided during the filling operation and during application. Fill the spray tank at least one-fourth full of clean water and while pump and agitator are running, add the recommended amount of BLADEX 4 WDS Herbicide to the tank, then fill with water to the desired level. For tank mixes, add the BLADEX 4 WDS Herbicide is completely suspended in the water, add the recommended amount of atrazine 4L, Sutan, Alachlor, Lasso, 4 EC, or Paraquat CL to the tank while filling with water to the desired level. For tank mixes with atrazine 80W, completely suspend the atrazine in water prior to adding BLADEX 4 WDS Herbicide. **Do not apply the BLADEX 4 WDS Herbicide/Paraquat in liquid fertilizer.**

Allow the pump to run several minutes before application to insure proper suspension and mixing of BLADEX 4 WDS Herbicide in the various mixtures. If BLADEX 4 WDS Herbicide or tank mix combinations of BLADEX 4 WDS Herbicide with atrazine are allowed to stand in a spray tank or hose tank for an extended period, several hours or overnight, provide good agitation for several minutes before application to insure proper suspension and mixing. Tank mix combinations of BLADEX 4 WDS Herbicide with Alachlor, Lasso, Sutan, or Paraquat should not be left in the spray tank for prolonged periods; batches of these should be mixed and applied the same day.

Apply the spray solution containing BLADEX 4 WDS Herbicide with the various mixtures evenly to the soil surface using 25-30 gallons per acre for corn and sorghum. BLADEX 4 WDS Herbicide may be applied by air using a minimum of 4 gallons of spray per acre.

Consult your local Agricultural Extension Agent for the proper determination of soil organic matter content and the most appropriate herbicide combination for your local conditions.

PREEMERGENCE

BLADEX 4 WDS Herbicide/Atrazine 80 W or 4L

Use BLADEX 4 WDS Herbicide + atrazine 80 W or 4L as tank mix combinations at a total rate of 1.0 to 1.4 pounds of total active ingredient per acre in corn and sorghum. For band applications apply proportionately less. Apply before planting, at planting, or after planting. Avoid removal of treated soil from seedbed during the planting operation. Adjust the total combined rate of application for a given soil class as possible according to the optimum rate given in the Preemergence Broadcast Application Rate Table for BLADEX 4 WDS Herbicide. Use two parts BLADEX 4 WDS Herbicide to one part atrazine in equal parts of both herbicides at a combined rate equal to the BLADEX 4 WDS Herbicide rate of application for a particular soil given in the table. The two parts BLADEX 4 WDS Herbicide to one part atrazine is suggested for grassier conditions. Suggested rates for these two ratios will be found below. Plant only corn or sorghum the year following the use of BLADEX 4 WDS Herbicide/ atrazine tank mixes.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 80 W (2 Parts BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Sand	DO NOT USE					
Light Sand		0.8 - 1.0	1.0 - 1.2	1.2 - 1.4	1.4 - 1.6	1.6 - 1.8
Sandy Clay	1.8 - 2.5	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8	2.8 - 3.0
Light Silty Clay	1.7 - 1.9	1.9 - 2.1	2.1 - 2.3	2.3 - 2.5	2.5 - 2.7	2.7 - 2.9
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Silty Clay	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8
Peat and Muck	NOT RECOMMENDED					

*If organic matter content between those listed, adjust the rate proportionately.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 80 W (1 Part BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Sand	DO NOT USE					
Light Sand		0.8 - 1.0	1.0 - 1.2	1.2 - 1.4	1.4 - 1.6	1.6 - 1.8

CHART CONTINUED IN NEXT COLUMN

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Sand	DO NOT USE					
Light Sand		0.8 - 1.0	1.0 - 1.2	1.2 - 1.4	1.4 - 1.6	1.6 - 1.8
Sandy Clay	1.8 - 2.5	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8	2.8 - 3.0
Light Silty Clay	1.7 - 1.9	1.9 - 2.1	2.1 - 2.3	2.3 - 2.5	2.5 - 2.7	2.7 - 2.9
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Silty Clay	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8
Peat and Muck	NOT RECOMMENDED					

*If organic matter content between those listed, adjust the rate proportionately.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 4L (2 Parts BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Sand	DO NOT USE					
Light Sand		0.8 - 1.0	1.0 - 1.2	1.2 - 1.4	1.4 - 1.6	1.6 - 1.8
Sandy Clay	1.8 - 2.5	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8	2.8 - 3.0
Light Silty Clay	1.7 - 1.9	1.9 - 2.1	2.1 - 2.3	2.3 - 2.5	2.5 - 2.7	2.7 - 2.9
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Silty Clay	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8
Peat and Muck	NOT RECOMMENDED					

*If organic matter content between those listed, adjust the rate proportionately.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 4L (1 Part BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Sand	DO NOT USE					
Light Sand		0.8 - 1.0	1.0 - 1.2	1.2 - 1.4	1.4 - 1.6	1.6 - 1.8
Sandy Clay	1.8 - 2.5	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8	2.8 - 3.0
Light Silty Clay	1.7 - 1.9	1.9 - 2.1	2.1 - 2.3	2.3 - 2.5	2.5 - 2.7	2.7 - 2.9
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Clay Loam	1.4 - 1.6	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6
Sandy Silty Clay	1.6 - 1.8	1.8 - 2.0	2.0 - 2.2	2.2 - 2.4	2.4 - 2.6	2.6 - 2.8
Peat and Muck	NOT RECOMMENDED					

*If organic matter content between those listed, adjust the rate proportionately.

BLADEX 4 WDS Herbicide/Alachlor (Lasso) 4 EC

Apply BLADEX 4 WDS Herbicide as a combination tank mix with Alachlor 4 EC herbicide immediately before or at planting or after planting, but before the crop has emerged. Avoid removal of treated soil from seedbed prior to or during the planting operation. When used with Alachlor 4 EC, BLADEX 4 WDS Herbicide should be used at 1/2 the rate shown on the Preemergence Broadcast Application Rates for BLADEX 4 WDS Herbicide. Sutan and Paraquat should be used at 1/2 the rate shown on the Preemergence Broadcast Application Rates for BLADEX 4 WDS Herbicide. Do not use BLADEX 4 WDS Herbicide in tank mixes with Alachlor 4 EC herbicide in the same application.

ALL C LABEL

LABEL FOR USE OF BLADEX® 4 WDS HERBICIDE IN TANK MIXES WITH ATRAZINE, ALACHLOR (LASSO), SUTAN+ OR PARAQUAT*

(Refer to label on container for ingredients, precautions, notice of warranty and use directions for BLADEX® alone and in these tank mixes.)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Clay	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W	Quarts of BLADEX 4 WDS + Pounds of Atrazine 80 W
Silt						
Silt Loam						
Sand						
Sand Loam						
Sandy Clay						
Sandy Clay Loam						
Peaty Muck	NOT RECOMMENDED					

*The rate is proportional to the percent organic matter between the levels; adjust the rate proportionately.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 4L (2 Parts BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Clay	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L
Silt						
Silt Loam						
Sand						
Sand Loam						
Sandy Clay						
Sandy Clay Loam						
Peaty Muck	NOT RECOMMENDED					

*The rate is proportional to the percent organic matter between the levels; adjust the rate proportionately.

BLADEX 4 WDS HERBICIDE BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND ATRAZINE 4L (1 Part BLADEX: 1 Part Atrazine)

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Clay	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L	Quarts of BLADEX 4 WDS + Quarts of Atrazine 4L
Silt						
Silt Loam						
Sand						
Sand Loam						
Sandy Clay						
Sandy Clay Loam						
Peaty Muck	NOT RECOMMENDED					

*The rate is proportional to the percent organic matter between the levels; adjust the rate proportionately.

BLADEX 4 WDS Herbicide: Alachlor (Lasso) 4 EC

Apply BLADEX 4 WDS Herbicide in a tank mix with Alachlor (Lasso) 4 EC herbicide to control weeds before planting or shortly after planting, but before the crop has emerged. Apply tank mix prior to or during the planting operation. When combined with Alachlor (Lasso) 4 EC herbicide, BLADEX 4 WDS Herbicide should be used at the rate shown in the Pre-emergence Broadcast Application Rates for BLADEX 4 WDS Herbicide Table. Do not use at 2 quarts per acre or greater in any soil type. Do not use in any soil type where the organic matter is less than 1%. Do not use in any soil type where the organic matter is 1% to 4%.

RECOMMENDED RATES PER ACRE FOR BLADEX 4 WDS HERBICIDE AND ALACHLOR (LASSO) 4 EC HERBICIDE MIXTURES BY SOIL TYPE

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil*					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Clay	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC	Quarts of BLADEX 4 WDS + Quarts of Alachlor 4 EC
Silt						
Silt Loam						
Sand						
Sand Loam						
Sandy Clay						
Sandy Clay Loam						
Peaty Muck	NOT RECOMMENDED					

*The rate is proportional to the percent organic matter between the levels; adjust the rate proportionately.

BLADEX 4 WDS Herbicide: Sutan + 6E

Use BLADEX 4 WDS Herbicide in a tank mix with Sutan + 6E herbicide in either a tank mix combination and incorporate immediately or apply BLADEX pre-emergence to control weeds before planting or shortly after planting. Do not use the rate recommended in the Broadcast Application Rates of Tank Mix Combinations of BLADEX 4 WDS Herbicide Table for application in any soil type. For incorporation, use power-driven cultivation equipment set for two to three inch depth in a tank mix combination of 2 quarts of BLADEX 4 WDS Herbicide plus 2 quarts of Sutan + 6E herbicide per acre. For thorough mixing, disc in two directions and follow with a harrow or roller. Do not use in any soil type where the organic matter is less than 1%. Do not use in any soil type where the organic matter is 1% to 4%.

ADDITIONAL WEEDS CONTROLLED BY BLADEX 4 WDS HERBICIDE PLUS SUTAN + 6E COMBINATIONS

- | | |
|--|---|
| GRASSES: Sandbar, Shattercane, Wild Cane, Texas Panhandle | SEDGES: Yellow nutsedge, Yellow nutgrass, Purple nutsedge, Purple nutgrass |
|--|---|

BROADCAST APPLICATION RATES PER ACRE OF TANK-MIX COMBINATIONS OF BLADEX 4 WDS AND SUTAN + 6E*

SOIL TYPE DESCRIPTION	Percent Organic Matter in Soil**					
	Less Than 1%	1%	2%	3%	4%	Over 4%
Clay	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E	Quarts of BLADEX 4 WDS + Quarts of SUTAN + 6E
Silt						
Silt Loam						
Sand						
Sand Loam						
Sandy Clay						
Sandy Clay Loam						
Peaty Muck	NOT RECOMMENDED					

**The rate is proportional to the percent organic matter between the levels; adjust the rate proportionately.

Do not use in any soil type where the organic matter is less than 1%. Do not use in any soil type where the organic matter is 1% to 4%. Do not use in any soil type where the organic matter is 4% to 6%. Do not use in any soil type where the organic matter is 6% to 8%. Do not use in any soil type where the organic matter is 8% to 10%. Do not use in any soil type where the organic matter is 10% to 12%. Do not use in any soil type where the organic matter is 12% to 14%. Do not use in any soil type where the organic matter is 14% to 16%. Do not use in any soil type where the organic matter is 16% to 18%. Do not use in any soil type where the organic matter is 18% to 20%. Do not use in any soil type where the organic matter is 20% to 22%. Do not use in any soil type where the organic matter is 22% to 24%. Do not use in any soil type where the organic matter is 24% to 26%. Do not use in any soil type where the organic matter is 26% to 28%. Do not use in any soil type where the organic matter is 28% to 30%. Do not use in any soil type where the organic matter is 30% to 32%. Do not use in any soil type where the organic matter is 32% to 34%. Do not use in any soil type where the organic matter is 34% to 36%. Do not use in any soil type where the organic matter is 36% to 38%. Do not use in any soil type where the organic matter is 38% to 40%. Do not use in any soil type where the organic matter is 40% to 42%. Do not use in any soil type where the organic matter is 42% to 44%. Do not use in any soil type where the organic matter is 44% to 46%. Do not use in any soil type where the organic matter is 46% to 48%. Do not use in any soil type where the organic matter is 48% to 50%.

BLADEX 4 WDS Herbicide: Paraquat CL

Use BLADEX 4 WDS Herbicide Paraquat CL in a tank mix combination in broadcast treatments for the kill of existing vegetation and residual control of weeds where corn will be planted directly into a cover crop established under or into previous crop residues. For maximum till or no-till corn production, use BLADEX 4 WDS Herbicide at the rate shown in the Pre-emergence Broadcast Application Rates for BLADEX 4 WDS Herbicide Table. Do not use in any soil type where the organic matter is less than 1%. Do not use in any soil type where the organic matter is 1% to 4%. Do not use in any soil type where the organic matter is 4% to 6%. Do not use in any soil type where the organic matter is 6% to 8%. Do not use in any soil type where the organic matter is 8% to 10%. Do not use in any soil type where the organic matter is 10% to 12%. Do not use in any soil type where the organic matter is 12% to 14%. Do not use in any soil type where the organic matter is 14% to 16%. Do not use in any soil type where the organic matter is 16% to 18%. Do not use in any soil type where the organic matter is 18% to 20%. Do not use in any soil type where the organic matter is 20% to 22%. Do not use in any soil type where the organic matter is 22% to 24%. Do not use in any soil type where the organic matter is 24% to 26%. Do not use in any soil type where the organic matter is 26% to 28%. Do not use in any soil type where the organic matter is 28% to 30%. Do not use in any soil type where the organic matter is 30% to 32%. Do not use in any soil type where the organic matter is 32% to 34%. Do not use in any soil type where the organic matter is 34% to 36%. Do not use in any soil type where the organic matter is 36% to 38%. Do not use in any soil type where the organic matter is 38% to 40%. Do not use in any soil type where the organic matter is 40% to 42%. Do not use in any soil type where the organic matter is 42% to 44%. Do not use in any soil type where the organic matter is 44% to 46%. Do not use in any soil type where the organic matter is 46% to 48%. Do not use in any soil type where the organic matter is 48% to 50%.

EPA Reg. No. 201-281-AA

SHELL CHEMICAL COMPANY, A Division of Shell Oil Company, AGRICULTURAL CHEMICALS, SAN RAMON, CA 94583

ALL COPY UNLESS OTHERWISE INDICATED IS SOLID BLACK
LABEL SHOWN IS 1/2 SIZE