



ACCEPTED
 Oct 13, 1970
 UNDER THE FEDERAL INSECTICIDE ACT
 FOR EMISSIONS CONTROL REGISTERED
 ED UNDER NO. 24-288 SUBJECT
 TO APPROVED COMMENTS.

GARDONA[®] 2 INSECTICIDE EMULSIBLE CONCENTRATE

Contains 2 lb. GARDONA Insecticide per gallon

WARNING! KEEP OUT OF REACH OF CHILDREN

GENERAL DIRECTIONS

GARDONA[®] Insecticide 2 E.C. is suitable for use in conventional hydraulic sprayers. Time applications in accordance with State and Federal recommendations; consult the State Agricultural Extension Service.

USE AND DIRECTIONS FOR APPLICATION

GARDONA 2 E.C. is for use on corn (sweet corn, popcorn, field corn) for the control of corn earworm, fall armyworm, and European corn borer.

For the control of corn earworm and fall armyworm feeding in the whorls prior to the development of tassels or silk use 3 to 4 pints GARDONA 2 E.C. per acre making application directly over the plants so that the spray will run into the whorls. Treatment should begin as soon as damaging numbers are apparent. Spray weekly or once every 3 to 4 days, depending on infestation pressure from the time the plant emerges until silking.

To protect ears from earworm larvae use 3 to 6 pints GARDONA 2 E.C. per acre. Direct the spray at the ear zone to insure uniform coverage of silks. Begin application on the day silks appear and continue until silks are dry and no longer attractive to moths for egg deposition. Under mid-season growing conditions and moderate to heavy infestation pressure, application intervals of 1 to 2 days may be necessary. During periods of slower growth and lighter infestation pressure, application intervals may be lengthened to 3 to 5 days. For maximum effectiveness spray should be directed towards the ear zone to thoroughly cover ears and silks and timed with larvae emergence.

GARDONA 2 E.C. can be applied up to the day of harvest for corn grown for human con-

ACTIVE INGREDIENTS	BY WEIGHT
2-Chloro-1-(2,4,5 trichlorophenyl) vinyl dimethyl phosphate	24.3%
Xylene	60.9%
INERT INGREDIENTS	14.8%
	TOTAL 100.0%

USDA Reg. No. 201-238

Net Contents: 30 GALLONS

sumption. Treated corn forage can be fed to cattle providing a five day interval exists between final treatment and day of feeding.

For first-brood European corn borer larvae ground application use 3 to 4 pints GARDONA 2 E.C. per acre in 10 to 15 gallons of water. Use two to three nozzles directly over the row. For second-brood European borer larvae use four nozzles per row (two on each side) directed towards the ear zone. Consult your local State Agricultural Extension Service for timing of applications.

WARNING!

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Harmful if swallowed. Apply this product only as specified on this label.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

SHELL CHEMICAL COMPANY, A DIVISION OF SHELL OIL COMPANY, AGRICULTURAL CHEMICALS DIVISION, NEW YORK, N.Y. 10020