[201-292]

ACCEPTED [illegible]
[illegible] INSECTICIDE
[illegible] RODENTICIDE
[illegible] POISON REGISTER
[illegible] NO. [illegible] SUBJECT
[illegible] ACHED COMMENTS.

PHOSDRIN® INSECTICIDE

EMULSIBLE CONCENTRATE NO. 2

(Contains 2.0 lbs. PHOSDRIN Insecticide Per Gal.)

ACTIVE INGREDIENTS	BY WEIGHT
alpha Isomer of 2-carbomethoxy-1 methylvinyl dimethyl phosphate	15.2%*
Related Compounds	10.2%*
Petroleum Hydrocarbons (Xylene)	68.6%
INERT INGREDIENTS	6.0%
TOTAL	100.0%

*Equivalent to 25.4% w PHOSDRIN Insecticide

USDA Reg. No. 201-292

DANGER! POISON

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS

SEE ANTIDOTE AND OTHER PRECAUTIONS BELOW

Read the Directions Carefully and Follow Them at all Times

USE AND APPLICATION DIRECTIONS

Shell PHOSDRIN Insecticide Emulsible Concentrate can be used in conventional hydraulic sprayers, low-volume ground applicators, or airplane sprayers. Use this product only for recommended purposes and at recommended dosages. For best results consult the State Agricultural Extension Service for exact dosage and timing recommendations.

Mix with water by vigorous agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

Apply when insects first appear and repeat application as often as necessary to maintain control. Apply in a suitable volume of water to give uniform coverage of foliage. For application by airplane, dilute this product with 3 to 10 gallons of water per acre.

Fruit and Nut Crops: Use recommended dosages in sufficient water to obtain thorough spray coverage, but do not exceed indicated maximum gallonage for that crop. If concentrate ground application is employed, do not apply less than 15 gallons of spray per acre, and adjust dosage of PHOSDRIN Insecticide Emulsible Concentrate to obtain same dosage per acre as recommended for full coverage application.

Field and Vegetable Crops: Use 10-125 gallons of spray per acre when using ground application equipment.

Avoid drift of this product to crops for which it is not registered or to crops for which its use may be registered but application would be closer to harvest than recommended.

* Consult your State Department of Agriculture or local Agricultural Extension Service for information on the time interval between application and safe re-entry to the treated field.

FRUIT AND NUT CROPS

APPLES, PEACHES, PEARS, PLUMS: For control of aphids and mites use [illegible] pint per 100 gals. water.

Do not exceed 1200 gallons of finished spray per acre on apples and pears when 1 pint is used. For control of grasshoppers, lygus bugs and red-banded leafroller use 1 to 1½ pts. per 100 gals. water. Do not exceed 1100 gallons of finished spray per acre on apples and pears when 1½ pts. is used. Do not treat within one day of harvest.

CHERRIES (SOUR ONLY): For control of aphids and mites use 1 to 1½ pts. per 100 gals. water. Do not treat within 2 days of harvest.

CITRUS (ORANGES, LEMONS, GRAPEFRUIT): For control of aphids use [illegible] per acre in 200 or more gals. water. For control of fruit tree [illegible], orange tortrix and omnivorous leafroller use [illegible] qts. per acre in 500 gals. water. For control of Western tussock moth larvae [illegible] and pink scavenger caterpillar [illegible] 1,000 gals. water. Allow at least 7 days between [illegible] within [illegible] day of harvest.

GRAPES: [illegible] aphids use 1 to [illegible] pts. per acre. Do not treat within [illegible]. For control of [illegible], leafhoppers, mites, red-banded leafrollers and [illegible] bugs use 1 to [illegible] qt. per acre. At [illegible] qt. [illegible] treat within 5 days of harvest.

RASPBERRIES: [illegible] water [illegible] 100 [illegible] water. Apply [illegible]

STRAWBERRIES: [illegible] water [illegible] grasshoppers [illegible] fruitfly [illegible] Do not treat within [illegible] day [illegible]

WALNUTS: [illegible] aphids [illegible]

FIELD CROPS

[illegible]

per acre. For control of cabbage looper, imported cabbage worm, false chinch bug, dipterous leafminer (adult), grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.

PEA VINES (FOR FORAGE ONLY): For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and cutworms (climbing) use 1 to 2 pts. per acre. Do not treat within one day of harvest.

SORGHUM (FOR FORAGE AND GRAIN): For control of corn earworm and webworms use 1 to 2 pts. per acre. For control of fall armyworms use 1 qt. per acre. Do not treat within 3 days of harvest.

VEGETABLE CROPS

ARTICHOKES: For control of aphids use ½ to 1 pt. per acre. For control of plume moth use 1 to 2 qts. per acre. Do not treat within 2 days of harvest.

BEANS: For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and Mexican bean beetle use 1 to 2 pts. per acre. Do not treat within one day of harvest.

BEETS (INCLUDING TOPS): For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, cutworm (climbing), dipterous leafminer (adult), false chinch bug, grasshoppers, imported cabbage worm, leafhopper, mites and salt marsh caterpillar use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.

BROCCOLI, CABBAGE: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1 to 2 pts. per acre. Do not treat within 1 day of harvest. For hard to kill insects, including aphids, use 2 qts. per acre but do not treat within 3 days of harvest.

BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, KALE: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt marsh caterpillar, mites, cutworm (climbing), dipterous leafminer (adult) and lygus bugs use 1 to 2 pts. per acre. Do not treat within 3 days of harvest. For hard to kill insects, including aphids, use 2 qts. per acre, but do not treat within 7 days of harvest.

CARROTS: For control of aphids use ½ to 1 pt. per acre. For control of leafhoppers, lygus bugs, mites, cabbage looper, dipterous leafminer (adults), cutworms (climbing) and salt marsh caterpillar use 1 to 2 pts. per acre. Do not treat within [illegible] days of harvest.

CELERY: For control of aphids use ½ to 1 pt. per acre. For control of dipterous leafminer (adults), lygus bug, salt marsh caterpillar, leafhoppers, cabbage loopers and mites use 1 to 2 pts. per acre. Do not treat within 3 days of harvest. For hard to kill insects including aphids use 2 qts. per acre but do not treat within [illegible] days of harvest.

CORN (FIELD, SWEET AND POPCORN, FOR GRAIN ONLY): For control of aphids use ½ to 2 pts. per acre. Do not treat within 1 day of harvest.

CUCUMBERS (OUTDOOR): For control of aphids [illegible]. For control of grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.

EGGPLANT, PEPPERS: For control of aphids [illegible]. For control of grasshoppers, leafhoppers and mites [illegible]. Do not treat within [illegible] days of harvest.

LETTUCE: For control of aphids [illegible] corn earworm, cutworms (climbing), dipterous leafminer [illegible], cabbage looper, imported cabbage worm, grasshoppers, mites, lygus bugs, salt marsh caterpillar, false chinch bug and thrips [illegible] per acre. Do not treat within [illegible] days of harvest. [illegible] but do not treat within [illegible] harvest.

MELONS (INCLUDING CANTALOUPE, HONEYDEW MELONS, MUSKMELONS), WATERMELONS: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, dipterous leafminer (adult), leafhoppers, mites, false chinch bug, salt marsh caterpillar [illegible] per acre. To control [illegible] cutworms [illegible]

[illegible]

worms (climbing) use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.

PEAS (INCLUDING VINES): For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and cutworms (climbing) use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.

POTATOES: For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 pt. per acre. Do not treat within 1 day of harvest.

SPINACH: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, dipterous leafminer (adults), cutworms (climbing), salt-marsh caterpillar and false chinch bug use 1 to 2 pts. per acre. Do not treat within 4 days of harvest. For hard to kill insects, including aphids, use 2 qts. per acre, but do not treat within 4 days of harvest.

SUMMER SQUASH: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, dipterous leafminer (adults), leafhoppers, lygus bugs, mites, false chinch bugs, salt marsh caterpillar, cutworms (climbing), grasshoppers use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.

TOMATOES (OUTDOOR): For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.

TURNIPS: For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, false chinch bug and dipterous leafminer (adults) use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.

GREENHOUSE

Diluted PHOSDRIN Insecticide Emulsible Concentrate should be applied in a closed greenhouse [illegible] and ventilators [illegible] entrances [illegible] post warning signs and take whatever precautions are necessary to prevent unprotected humans and domestic animals from entering the treated area. The operator must wear a full face mask of a type found adequate for mevinphos (PHOSDRIN Insecticide) protection.

Apply any time that the greenhouse ventilators can remain closed for [illegible] hours without endangering [illegible] from high temperatures, such as early morning, cloudy days or late afternoon. If application is made during the day, keep the greenhouse closed tightly for at least two hours, then ventilate the enclosure thoroughly for an hour before resuming regular work. If application is made in the late afternoon, the house can be kept closed all night. On the following morning, ventilate for one hour to render the air safe for regular work.

LETTUCE (INCLUDING LEAF, BIBB AND ROMAINE): For control of aphids, corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, lygus bugs, salt marsh caterpillars, false chinch bugs, thrips use 2½ pts. [illegible] per 50,000 [illegible] ft. Do not treat within 10 days of harvest.

DANGER

[illegible]

Wear [illegible] clean protective clothing and [illegible] Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type found adequate for mevinphos (PHOSDRIN Insecticide) protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 10 minutes [illegible] with soap and water [illegible] before re-use.

[illegible]

weather conditions favor drift from areas treated [illegible] runoff is likely to occur.

This product is poisonous to fish and wildlife. Bi[illegible] areas may be killed. Keep out of lakes, ponds and [illegible] in any manner not specified on the label.

DECONTAMINATE AND DESTROY USED CONTAINERS [illegible] purpose. Decontaminate containers by washing [illegible] strongly alkaline detergent solution and rinsing [illegible] of rinsings in a way as not to constitute a hazard [illegible] supplies.

Destroy containers by crushing or puncturing. [illegible] burned stay out of smoke.

Note: DANGER! Avoid skin and eye contamination [illegible] caustic solutions.

ANTIDOTE AND FIRST AID TR[illegible]

ATROPINE IS THE EMERGENCY ANTIDOTE FOR P[illegible] POISONING.

CALL A PHYSICIAN IN ALL CASES OF SUSPECTED [illegible]

INTERNAL: If the material has been swallowed, [illegible] mediately. This may be done by introducing a [illegible] or by giving warm salt water (1 tablespoon of salt [illegible]). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONS[illegible]

EXTERNAL: If the material has been spilled on [illegible] remove patient from the vicinity of the insectic[illegible] taminated clothing, and wash skin with soap and [illegible] material gets into the eyes, wash immediately w[illegible] at least ten minutes.

IF WARNING SYMPTOMS APPEAR: (See WARNING S[illegible]) patient prone and quiet. Start artificial respirati[illegible] tient is not breathing. Transport the patient imme[illegible] hospital.

NOTE TO PHYSIC[illegible]N

WARNING SYMPTOMS: Symptoms [illegible] weakness [illegible] chest, blurred vision, non-reactive pin-point pup[illegible] ing, nausea, vomiting, diarrhea and abdominal c[illegible]

TREATMENT: Atropine is the specific therapeutic [illegible] against parasympathetic nervous stimulation. If th[illegible] sympathetic stimulation, atropine sulfate [illegible] intravenously in doses of 2 to [illegible] milligrams [illegible]

Pralidoxime chloride (2-PAM) [illegible] be [illegible] antidote in addition to and with [illegible] maintaining [illegible] adults, an initial dose of 1 gram of 2-PAM should [illegible] intravenously in 250 cc of saline over 15 to 30 [illegible] not practical, 2-PAM may be administered slow[illegible] in a 5 percent solution in water over not l[illegible] After about an hour a second dose of 1 gram of 2-[illegible] if muscle weakness [illegible] For inf[illegible] dose of 2-PAM is [illegible]

[illegible]

[illegible] Oxygen [illegible] cholinesterase inhibitors may [illegible] without warning. [illegible] cholinestera[illegible] further exposure until [illegible] regeneration is determined by blood test.

NOTICE OF WARRANT[illegible]

SHELL CHEMICAL COMPANY MAKES NO WARRANTY [illegible] FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPR[illegible] cerning this product [illegible] extend be[illegible] [illegible] with [illegible]

NET CONTENTS: 30 G[illegible]

SHELL CHEMICAL COMPANY A Division of Shell Oil Company AGRICULTURAL DIVISION SAN RAMON, CA 94583

PHOSDRIN® INSECTICIDE

EMULSIBLE CONCENTRATE NO. 2

(Contains 2.0 lbs. PHOSDRIN Insecticide Per Gal.)

ACTIVE INGREDIENTS	BY WEIGHT
alpha Isomer of 2 carbomethoxy 1-methylvinyl dimethyl phosphate	[illegible]*
Related Compounds	[illegible]*
Petroleum Hydrocarbons (Xylene)	68.[illegible]
INERT INGREDIENTS	6.0%
TOTAL	100.0%

*Equivalent to 25.4%w PHOSDRIN Insecticide

USDA Reg. No. 201-292

DANGER! POISON

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS

SEE ANTIDOTE AND OTHER PRECAUTIONS BELOW

per acre. For control of cabbage looper, imported cabbage worm, false chinch bug, dipterous leafminer (adult), grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.
PEA VINES (FOR FORAGE ONLY) For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and cutworms (climbing) use 1 to 2 pts. per acre. Do not treat within one day of harvest.
SORGHUM (FOR FORAGE AND GRAIN) For control of corn earworm and webworms use 1 to 2 pts. per acre. For control of fall armyworms use 1 qt. per acre. Do not treat within 3 days of harvest.

VEGETABLE CROPS

ARTICHOKES: For control of aphids use ½ to 1 pt. per acre. For control of plume moth use 1 to 2 qts. per acre. Do not treat within 2 days of harvest.
BEANS For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and Mexican bean beetle use 1 to 2 pts. per acre. Do not treat within one day of harvest.
BEETS (INCLUDING TOPS). For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, cutworm (climbing), dipterous leafminer (adult), false chinch bug, grasshoppers, imported cabbage worm, leafhopper, mites and salt-marsh caterpillar use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.
BROCCOLI, CABBAGE For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1 to 2 pts. per acre. Do not treat within 1 day of harvest. For hard to kill insects including aphids, use 2 qts. per acre but do not treat within 3 days of harvest.
BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, KALE For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and lygus bugs use 1 to 2 pts. per acre. Do not treat within 3 days of harvest. For hard to kill insects including aphids use 2 qts. per acre but do not treat collards and kale within 7 days of harvest.
CARROTS For control of aphids use ½ to 1 pt. per acre. For control of leafhoppers, lygus bugs, mites, cabbage looper, dipterous leafminer (adult), cutworms (climbing) and salt-marsh caterpillar use 1 to 2 pts. per acre. Do not treat within [illegible] days of harvest.
CELERY For control of aphids use ½ to 1 pt. per acre. For control of dipterous leafminer (adults), lygus bug, salt-marsh caterpillar, leafhoppers, cabbage loopers and mites use 1 to 2 pts. per acre. Do not treat within 3 days of harvest. For hard to kill insects including aphids use 2 qts. per acre but do not treat within [illegible] days of harvest.
CORN (FIELD, SWEET AND POPCORN) FOR GRAIN ONLY For control of [illegible] use ½ to 2 pts. per acre. Do not treat within [illegible] of harvest.
CUCUMBERS (OUTDOOR) For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.
EGGPLANT, PEPPERS For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 2 days of harvest.
LETTUCE For control of aphids use ½ to 1 pt. per acre. For control of corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, lygus bugs, salt-marsh caterpillar, false chinch bug and thrips use 1 to 2 pts. per acre. Do not treat within [illegible] days of harvest. For hard to kill insects including aphids use 2 qts. per acre but do not treat within [illegible] days of harvest.
MELONS, INCLUDING CANTALOUPES, HONEY DEW MELONS, MUSKMELONS, WATERMELONS For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, dipterous leafminer (adult), leafhoppers, lygus bugs, mites, false chinch bugs, salt-marsh caterpillars, [illegible] and grasshoppers use 1 to 2 pts. per acre. To control rind worms, cabbage looper, [illegible] [illegible]

[illegible]

worms (climbing) use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.
PEAS (INCLUDING VINES) For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers, mites and cutworms (climbing) use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.
POTATOES For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 pt. per acre. Do not treat within 1 day of harvest.
SPINACH For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, dipterous leafminer (adults), cutworms (climbing), salt-marsh caterpillar and false chinch bug use 1 to 2 pts. per acre. Do not treat within 4 days of harvest. For hard to kill insects, including aphids use 2 qts. per acre, but do not treat within 4 days of harvest.
SUMMER SQUASH For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, dipterous leafminer (adults), leafhoppers, lygus bugs, mites, false chinch bugs, salt marsh caterpillar, cutworms (climbing), grasshoppers use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.
TOMATOES (OUTDOOR) For control of aphids use ½ to 1 pt. per acre. For control of grasshoppers, leafhoppers and mites use 1 to 2 pts. per acre. Do not treat within 1 day of harvest.
TURNIPS For control of aphids use ½ to 1 pt. per acre. For control of cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, false chinch bug and dipterous leafminer (adults) use 1 to 2 pts. per acre. Do not treat within 3 days of harvest.

GREENHOUSE

Diluted PHOSDRIN Insecticide Emulsible Concentrate should be applied in a closed greenhouse: close all doors, windows and ventilators. Lock or barricade all entrances, post warning signs, and take whatever precautions are necessary to prevent unprotected humans and domestic animals from entering the treated area. The operator must wear a full face mask of a type found adequate for mevinphos (PHOSDRIN Insecticide) protection.
Apply any time that the greenhouse ventilators can remain closed for two hours without endangering crops from high temperatures, such as early morning, cloudy days or late afternoons. If application is made during the day, keep the greenhouse closed tightly for at least two hours, then ventilate the enclosure thoroughly for an hour before resuming regular work. If application is made in the late afternoon, the house can be kept closed all night. On the following morning, ventilate for one hour to render the air safe for regular work.
LETTUCE (INCLUDING LEAF, BIBB AND ROMAINE) For control of aphids, corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, lygus bugs, salt-marsh caterpillars, false chinch bugs, thrips use 2⅔ pts. to 4⅔ pts. per 50,000 cu. ft. Do not treat within 10 days of harvest.

DANGER

[illegible]

Do not swallow. Do not get in eyes, on skin, or on clothing. Do not breathe vapor. Do not contaminate food or feed products. Keep away from [illegible]

Wear clean rubber gloves, clean protective clothing and goggles or face shield. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type found adequate for mevinphos (PHOSDRIN Insecticide) protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least 10 minutes and get medical attention. In case of contact remove contaminated clothing and wash skin with soap and water. Wash clothing before reuse. [illegible]

[illegible]

weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.
This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.
DECONTAMINATE AND DESTROY USED CONTAINERS Do not reuse for any purpose. Decontaminate containers by washing thoroughly with a strongly alkaline detergent solution and rinsing with water. Dispose of rinsings in a way as not to constitute a hazard or contaminate water supplies.
Destroy containers by crushing or puncturing, then bury, or burn. If burned stay out of smoke.
Note: DANGER! Avoid skin and eye contamination when handling strong caustic solutions.

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE IS THE EMERGENCY ANTIDOTE FOR PHOSDRIN INSECTICIDE POISONING.
CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.
INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.
EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.
IF WARNING SYMPTOMS APPEAR: (See WARNING SYMPTOMS below) Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.
TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred.
Pralidoxime chloride (2-PAM chloride) may also be used as an effective antidote in addition to and while maintaining full atropinization. In adults, an initial dose of 1 gram of 2-PAM should be injected, preferably as an infusion in 250 cc of saline over a 15 to 30 minute period. If this is not practical, 2-PAM may be administered slowly by intravenous injection as a 5 percent solution in water over not less than two minutes. After about an hour, a second dose of 1 gram of 2-PAM will be indicated if muscle weakness has not been relieved. For infants and children the dose of 2-PAM is 0.25 grams.
Morphine is an improper treatment.
Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by blood test.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its use, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

NET CONTENTS: 30 GALLONS

[illegible] CHEMICAL COMPANY, A Division of Shell Oil Company AGRICULTURAL DIVISION, SAN RAMON, CA 94583