

DILUTION TABLE

CONCENTRATION	QUANTITY OF WATER			
	1 Gallon	5 Gallons	25 Gallons	100 Gallons
Amount of Poultry Spray Emulsifiable for 0.50% Concentration	3 Ozs.	13 Ozs.	2 Quarts	2 Gallons
Amount of Poultry Spray Emulsifiable for 1.00% Concentration	6 Ozs.	26 Ozs.	4 Quarts	4 Gallons

NOTE: Agitate mixture thoroughly before application.

PRECAUTIONS IN USING

CAUTION! Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

**AGRICULTURAL DIVISION
HEALTH & NUTRITION
SAN RAMON, CA. 94583**

POULTRY S.
EMULSIFIABLE—
Contains **RABON®** Insecti

**Controls HOUSE FLIES, NORTHERN
CHICKEN MITES, and LICE—On or in
&
RESIDUAL FLY CONTROL and**

ACTIVE INGREDIENTS	BY WEIGHT
2-Chloro-1,2,4,5-trichloro- phenyl vinyl dimethyl phosphate	24.0
Petroleum distillates	61.4
INERT INGREDIENTS	14.6
TOTAL	100.0



MUST BE DILUTE

201-243
ACCEPTED
DECE 1982

DIRECTIONS FOR USE

TO CONTROL	TYPE HOUSING	STRENGTH TO USE	APPLICATION	REMARKS
LICE & MITES	Wire Cages	0.5%	1 gal. 100 birds	Apply directly to birds, spray vent and fluff areas from below. Repeat when necessary. Do not repeat more often than every 14 days. For individual bird treatment apply 1 ounce of the mixture per bird.
	NOTE: For maximum lasting control of the northern fowl mite, penetration of the feathers around the vent area is absolutely essential. Use power sprayer at 100-125 psi at NO LESS THAN RECOMMENDED RATE. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.			
	Floor Management Litter	0.5%	1-2 gal. 1000 sq. ft.	Apply evenly for penetration to litter surface. Also apply thoroughly to walls, roosts, cracks, crevices and interiors. Spray birds lightly.
	Roost Paint	1.0%	1 pt. 100 sq. ft.	Treat with brush or spray thoroughly, particularly cracks and crevices.
FOWL TICK (Blue Bug)	All Types	1.0%	1 gal. 100-150 sq. ft.	Apply thoroughly to walls, ceilings, floors, cracks and crevices with a power sprayer.
FLIES RESIDUAL	All Types	1.0%	1 gal. 500-1000 sq. ft.	Apply thoroughly to point of run-off to walls, ceilings and where flies congregate and feed. Do not apply to the birds.
FLIES LARVAE	Wire Cages	1.0%	1 gal. 100 sq. ft. of droppings	A thorough coverage and penetration is essential for initial control.
NOTE: Regardless of the condition of the droppings apply enough spray to penetrate and kill the larvae. Less spray will be required thereafter provided a regular schedule is followed.				

SHELL CHEMICAL COMPANY
A Division of Shell Oil Company

ACL-1456 A

LABEL SHOWN IS ACTUAL SIZE
GREY AREA IS SOLID YELLOW
ALL COPY UNLESS OTHERWISE INDICATED IS SOLID BLACK