


SHELL CHEMICAL COMPANY
A Division of Shell Oil Company
Agricultural Division

San Ramon, California



Shell Poultry Spray 505 Wettable Powder
(Contains DDT Insecticide)

Controls northern fowl mites, chicken mites, and lice
on or around chickens
and
Residual Fly Control and Larvicide

ACTIVE INGREDIENTS	PERCENT BY WEIGHT
2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate	50.0
INERT INGREDIENTS	50.0
TOTAL	100.0

USDA Reg. No. 231- 296

Net Contents _____

CAUTION: KEEP OUT OF REACH OF CHILDREN.

PRECAUTIONS IN USING

Harmful if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of food and feedstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EITHER OF IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

BEST DOCUMENT AVAILABLE

This label has been reviewed and approved for compliance in regulatory
[Signature]

Page 1 of 3

[Handwritten mark]

7

DIRECTIONS FOR USE

<u>TYPE HOUSING</u>	<u>STRENGTH TO USE</u>	<u>APPLICATION</u>	<u>REMARKS</u>
<u>LICE AND MITES FOR PDL</u>			
WIRE CAGES	0.5%	1 gal/1000 sq. ft.	Apply directly to birds, spray vent and fluff areas from below. Repeat when necessary. Do not repeat more often than every 14 days. For individual bird treatment apply 1 ounce of the mixture per bird.

NOTE: For maximum lasting control of the northern fowl mite penetration of the feathers around the vent area is absolutely essential. Use power sprayer at 100-125 psi at no less than recommended rate. More attention must be given each individual bird when using low-pressure equipment. Treat roosters carefully and thoroughly to avoid reinfestation in breeding flocks.

FLOOR MANAGEMENT

Litter	0.5%	1-2 gal/1000 sq. ft.	Apply evenly for penetration to litter surface. Also apply thoroughly to walls, roosts, cracks, crevices and interiors. Spray birds lightly.
Litter	50% WP	2.5 oz/100 sq. ft.	Treat evenly and thoroughly.
Lust Box	50% WP	2.5 oz/50 birds	Mix evenly throughout top layer of contents.
Roost Point	1.0%	1 pt/100 ft.	Treat with brush or sprayer thoroughly, particularly cracks and crevices.

INDIVIDUAL BIRD TREATMENT

ALL TYPES	1.0%	1 gal/900-1000 sq. ft.	Apply thoroughly to point of contact to walls, ceiling and where flies congregate and roost. Do not apply to the birds.
-----------	------	------------------------	---

POWL TICK (DELTACENTRUM) TREATMENT

ALL TYPES	1.0%	1 gal/1000-1500 sq. ft.	Apply thoroughly to walls, ceiling, cracks and crevices with a power sprayer.
-----------	------	-------------------------	---

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

<u>TYPE HOUSING</u>	<u>STRENGTH OF USE</u>	<u>APPLICATION</u>	<u>REMARKS</u>
<u>LARVICIDE</u>			
WIRE CAGES			
Initial Control	1.0%	1 gal/100 sq. ft. of droppings	A thorough coverage and penetration is essential for initial control.

NOTE: Regardless of the condition of the droppings apply enough spray to penetrate and kill the larvae. Less spray will be required thereafter provided a regular schedule is followed.

DILUTION TABLE

Shell Poultry Spray 4.4 W.P. in Water

<u>CONCENTRATION</u>	<u>1 GAL</u>	<u>5 GALS</u>	<u>25 GALS</u>	<u>100 GALS</u>
0.5%	2.5 tbs.	6.5 oz.	2 lb.	8 lb.
1.0%	5 tbs.	13 oz.	4 lb.	16 lb.

NOTE: Agitate mixture thoroughly before application.

BEST DOCUMENT AVAILABLE