



POULTRY INSECTICIDE

MY-T-CIDE* Poultry Spray Emulsifiable

(Contains 2.0 lb. VAPONA® INSECTICIDE per gallon)
Controls: Northern Fowl Mite, Chicken Red Mite and Lice on or around Chickens



DANGER!  **POISON**

KEEP OUT OF THE REACH OF CHILDREN
AND DOMESTIC ANIMALS.
(See Antidote and other Precautions on label)

ACTIVE INGREDIENTS	BY WEIGHT
2,2-Dichlorovinyl dimethyl phosphate	21.8%*
Related Compounds	1.6%*
Petroleum Hydrocarbons	65.6%
INERT INGREDIENTS	11.0%
TOTAL	100.0%

*Equivalent to 23.4% VAPONA® INSECTICIDE

EPA Reg. No. 201-302

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for VAPONA® Insecticide poisoning. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR:

Keep patient prone and quiet.

Start artificial respiration immediately if patient is not breathing.

Transport the patient immediately to the nearest hospital.

NOTICE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

(continued in column at right)

DIRECTIONS FOR USE

NORTHERN FOWL MITES, CHICKEN RED MITES AND LICE

Type Housing	Application	Remarks
Wire Cages	1 gal./100 birds	Apply the 0.5% spray with high pressure equipment (100 psi or more) from below to the vent and fluff area. Do not treat more often than every 14 days. Do not contaminate feed or water.
Floor Litter, Roosts and Nests	1½ Quarts/ 100 sq. ft.	Apply the 0.5% spray evenly to litter surface. Retreat 10-14 days later. Do not wet nests excessively. Do not contaminate feed or water.

FLY CONTROL

All Types	1 Quart/1000 sq. ft.	Apply the 0.5% spray as a coarse wet spray to walls, beams, rafters, window sills and where flies congregate.
	1 Quart/8000 cu. ft. (as a fog)	Mix 1 pint of MY-T-CIDE with 7 gallons of fuel oil and apply recommended amount with any suitable fogger or mister. Do not contaminate feed or water.

LARVICIDE

Upper Caged Layers and on piled manure	2 Quarts/100 sq. ft. of manure	Apply the 0.5% spray thoroughly as a coarse spray to the manure.
--	--------------------------------	--

DILUTION TABLE

Preparation of 0.5% spray

Amount of My-T-Cide	Amount of Water
1 Pint	6 Gal.
1 Quart	12½ Gal.
1 Gallon	50 Gal.

Net Contents: 5 Gal.

Morphine is contraindicated.

Clear chest by postural drainage.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

PRECAUTIONS IN USING

Shell Poultry Insecticide MY-T-CIDE Poultry Spray Emulsifiable and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See Antidote and warnings statement.)

Store MY-T-CIDE Poultry Spray in a safe place. Children or animals must not be able to reach or handle MY-T-CIDE Poultry Spray or diluted emulsion. It is recommended that only the amount of diluted emulsion that will be used immediately be prepared from MY-T-CIDE Poultry Spray.

Mix MY-T-CIDE Poultry Spray into water outdoors or in a well ventilated room. Wash off the outside of the bottle of MY-T-CIDE Poultry Spray in running water before storing away.

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on label.

Do not contaminate feed, water or foodstuffs, milk or milking utensils.

DANGER

Shell Poultry Insecticide MY-T-CIDE Poultry Spray Emulsifiable is poisonous if swallowed, inhaled or absorbed through skin or eyes. Do not get on skin or eyes. Do not breathe fumes.

Avoid exposure to fumes. Wear clean rubber gloves, protective clothing and goggles when handling MY-T-CIDE Poultry Spray to prepare emulsions. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for DDVP (VAPONA Insecticide) protection. Keep all unprotected persons out of operational area. Not for use or storage in or around the home.

In case of contact, wash immediately with soap and water. Wash thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

DECONTAMINATE AND DESTROY USED CONTAINERS.

Decontaminate containers by washing thoroughly with a strong solution of alkali and detergent and rinsing with water.

Destroy containers by crushing or puncturing; then bury or burn. If burned, stay out of smoke.

Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning the product or its use, which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

SHELL CHEMICAL COMPANY, A Division of Shell Oil Company, AGRICULTURAL DIVISION, HEALTH & NUTRITION, SAN RAMON, CA 94583

*Shell Trademark

ACL-1608B