### GENERAL DIRECTIONS


RABON® Livestock Dust is a ready-to-use insecticide dust which can be applied by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter.

### USE DIRECTIONS

USE ORKEUTIONS		
AMMAL.	INSECT	REMARKS
Beef Cattle	Horn Flies, Face Flies & Lice	HAND BUSTING — Apply approximately 2 oz. of dust by shake can, retary duster or by spoon to the upper portions of the back, nack and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.  SELF-TREATING DUST BAG—  FUNCES USE—Put RABON® Livestock Dust in cotton cloth or double buriap bags or use pre-packed weather-proof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or buriap bags from weather.  FREE CHOICE USE—Use the same dust bags as above but place in leading sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The free choice aids in the control of lice.
Swine	Lice	WARE DUSTING—Apply 3-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more ethan than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One its of 3% dust should be applied per 150 sq. ft. o bedding.

## RABON° LIVESTOCK DUST A Ready-To-Use Insecticide Dust

CONTROLS HORN FLIES & LICE & DAIRY CATTLE and CONTROLS LICE ON SHIRE





NET WT.: 121/2 LBS.




JAN' 21

### RABON' LIVESTOCK DUST

A Ready-Te-Use insecticide Dust

CONTROLS HORN FLIES & LICE & aids in control of FACE FLIES ON DEEF & DAIRY CATTLE and COUTROLS LICE ON SAME





NET WT .: 12 1/2 LBS.



### PRECAUTIONS IN USING

CAUTION: Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

### NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR MIPLIED, concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this tabel.

SHELL CHEMICAL COMPANY
A Division of Shell Oil Company
ANIMAL HEALTH
SAN RAMON, CALIFORNIA 94583

**ACD-2648** 

applied by hand or from last application

z. of dust by shake per portions of the n aid in the comtrol dust beneath the


weather-proof cator alleyways leadal blocks, or water rom weather

bags as above but edlots, near waterher. The free choice

conventional hand tial attention given aftention given an necessary but not severe infestations be treated. One the ft. of bedding

# RABON<sup>®</sup> LIVESTOCK DUST

CONTROLS HORN FLIES & LICE & aids in control of FACE FLIES ON BEEF & DAIRY CATTLE and CONTROLS LICE ON SWIME



TOT: :00.00°

### CAUTION

KEEP OUT OF REACH OF CHILDREN.

SEE CTHER CAUTIONS ON SIDE FANEL

EPA REG. NO. 201-319 AA

EPA EST 10543-MO-01

TAKE TIME





CAUCION: Harmful if swan skin, or an elothing, of water for 15 minutes aughty with suap and wavoid contamination of open flame. On not appoint aminate water by other product andy as spand or discard in a safe plant of discard in a safe plant.

FITNESS FOR ANY PURI teraing this product of the under normal condilabel

> SHEL A Divi

SAN R