

GENERAL DIRECTIONS

Shell Horse SPRAY-N-WIPE is a ready-to-use combination of insecticides and repellents that can be applied either as a wipe-on or spray. Prior to application, brush away dirt and dust from the animal's hair coat. To apply as a WIPE-ON, pour a small amount of insecticide repellent evenly onto a piece of cloth or insecticide applicator or mist and wipe on the hair coat, first against the hair and then with the hair. Apply from 1-2 oz. of material per animal As a SPRAY, pour the insecticide repellent into a spray applicator and apply from 1-2 fluid oz. of solution as a fine mist to the areas of the horse's body needing insect protection.

NOTE: It is not necessary to exceed 2 fluid oz. per mature horse, slightly less for younger animals and ponies. It may be desirable to repeat the application after the horse has been ridden hard or because of weather conditions. For insect protection around the horse's head, the WIPE-ON method is preferred.

-USE DIRECTIONS

INSECT PROBLEM	REMARKS
FACE FLIES HOUSE FLIES	Apply as a wipe-on, preferably, to the head area. Pay particular attention to around the nostrils and eyes but avoid direct contact with the eye itself.
STABLE FLIES	Apply as a wipe-on or spray to the legs and flanks, leaving no unprotected area. More frequent applications may be necessary if the legs are exposed to high grass or water.
BITING GNATS	Apply as a wipe-on, preferably, to the head, neck, belly, and forelegs. Be sure to apply the insecticide repellent to the inside surfaces of the ears.
MOSQUITOES HORSE FLIES HOUSE FLIES DEER FLIES HORN FLIES	Apply as a wipe-on or spray to evenly cover the flanks, belly, and back of the horse or pony.

SOLIO BLUE

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. For use on horses only. Wash hands after use. Do not use, pour, spill or store near heat or open flame. Do not reuse container. Destroy when empty.

ACTIVE INGREDIENTS: (By Weight) 2-chioro-1-(2.4.5-trichlorophenyl)vinyl dimethyl phosphate* 1.00%; Pyrethrins 0.09%; Technical piperonyl butoxide 0.18% (consists of 0.14% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% related compounds); N-octyl bicycloheptene dicarboximide 0.30%; Di-n-propyl isocinchomeronate 0.50%; 2,3:4,5-bis (2-butylene) tetrahydro-2-furaldehyde 0.50%; Citronella 0il 0.1%; Petroleum distillates 96.27%. TOTAL 100%.

INERT INGREDIENTS: 1.06°ow.

RARON Insecticide.

EPA Reg. No. 201-339 AA

OBSERVE LABEL

TAKE TIME

DIRECTIONS

Made in U.S./

SOLID RED

ACD 299A

LABEL SHOWN IS ACTUAL SIZE

ALL COPY UNLESS OTHERWISE INDICATED IS SOLID BLACK

Transfer in the Arta Arta