



TECHNICAL NUDRIN* INSECTICIDE

FOR MANUFACTURING PURPOSES ONLY

AN INSECTICIDE FOR FORMULATING USE



ACTIVE INGREDIENTS	BY WEIGHT
S-methyl N-[(methylcarbamoyl)oxy]thioacetimidate	98%
INERT INGREDIENTS	2%
TOTAL	100%

EPA Reg. No. 201-353

PRECAUTIONS IN USING

Wear protective clothing, goggles and a mask or respirator suitable for protection against insecticide dust or spray mist, such as Willson Products "Agri-Tox-Z" R-15 Felt R-28 Cartiridge or Mines Safety Appliances Type H "Comfo." Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Do not contaminate or store near feed or foodstuffs.

DANGER! POISONOUS IF SWALLOWED OR INHALED

Do not breathe dust or spray mist.

Do not get in eyes, on skin, on clothing.

ANTIDOTE AND FIRST AID

Atropine is the emergency antidote for Shell NUDRIN Methomyl Insecticide poisoning. Call a physician at once in all cases of suspected poisoning.

INTERNAL: If swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. (NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.)

EXTERNAL: In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention; remove and wash contaminated clothing before re-use.

If warning symptoms appear (see warning symptoms below) keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest hospital.

DANGER  **POISON**

DANGER! KEEP OUT OF REACH OF CHILDREN.

See antidote and precautions statement.

Additional information pertinent to the needs of the formulator may be found in Technical Data Bulletin ADP:73-120.

NOTE TO PHYSICIAN

Warning Symptoms—Methomyl poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

Treatment—Atropine sulfate is the therapeutic antagonist of choice. Administer repeated doses, 1.2 to 2.0 mg intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured.

Do not use morphine or 2-PAM.

IMPORTANT—Not for use or storage in or around the home. Keep animals and unprotected persons out of operational areas during treatment and while there is danger of drift. Do not re-use container; bury when empty. Do not contaminate water when cleaning equipment or disposing of wastes or containers.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OR MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning the product or its uses which extend beyond the use of the product under normal conditions of use with the statements made on this label.

NET WEIGHT: 200 POUNDS

SHELL CHEMICAL COMPANY, A Division of Shell Oil Company, AGRICULTURAL DIVISION, SAN RAMON, CA. 94583

*Shell Trademark