DIRECTIONS: Place Flea Collar around dog's neck, adjust for proper fit and buckle in place. The collar must be worn loosely and in addition to the regular collar. If collar is worn too tightly, it may produce neck irritation. Cut off and dispose of the excess length. Flea kill begins at once and will continue for 3 months while the collar is worn. Ticks are harder to kill but those on the neck area will soon be killed and easily removed. Collar will continue to kill ticks for 8 weeks. For continuous protection, replace collar every 3 months.

Flea Collar can be worn with regular collar. Remove collar when washing dogreplace when dry.

CAUTION: Do not open protective pouch until ready to use. Do not allow children to play with this collar. Collar is intended for use only as an insecticide generator (Shell's U. S. Patents 2,956,073. 3.116,201 and 3.318,769) and is not to be taken internally by man or animals. Do not use on cats, whippets and greyhounds. Some dogs may be sensitive to this collar. Remove collar if there is evidence of irritation. Do not use on sick or convalescing dogs. Do not use any other pesticide on dog while collar is worn.

> Manufactured by SHELL CHEMICAL COMPANY A Division of Shell Oil Company AGRICULTURAL DIVISION **SAN RAMON, CA. 94583**

ACL-1694

EPA REG. NO. 201-356

201 - 2 - 2



FOR LONG-LASTING FLEA CONTROL AIDS IN TICK CONTROL-ESPECIALLY IN NECK AREA

CAUTION: Do not allow children to play with this collar — see cautions on back panel. ACTIVE INGREDIENTS BY WEIGHT

 2.2-dichlorovinyl dimethyl phosphate
 8.9%*

 Related Compounds
 0.7%*

 INERT INGREDIENTS
 90.4%

Equivalent to 9.6% w VAPONA Insecticide. TOTAL 100.0%

CONTENTS: 1 FLEA COLLAR

NET WEIGHT: 0.8 OZ.

LABEL SHOWN IS ACTUAL SIZE ALL COPY UNLESS OTHERWISE INDICATED IS SOLID BLACK

PED