DIRECTIONS: Place Flea Collar around cat's neck, adjust for proper fit and buckle in place. The collar must be worn loosely and in addition to the regular collar. If collar is worn too tightly it may produce neck irritation. Cut off and dispose of the excess length. Flea kill begins at once and will continue for 3 months while the collar is worn. For continuous protection, replace collar when effectiveness diminishes. When bathing cat, remove collar until cat is dry.

CAUTION: Do not open protective pouch until ready to use. Do not allow children to play with this collar. Do not use on Persians. Do not use on sick or convalescing cats. Remove collar if there is evidence of irritation. Do not use any other pesticides on cat while collar is worn. Collar is intended for use only as an insecticide generator (Shell's U. S. Patents 2.956.073, 3.116.201 and 3.318.769) and is not to be taken internally by man or animals.

> Manufactured by SHELL CHEMICAL COMPANY A Division of Shell Oil Company AGRICULTURAL DIVISION SAN RAMON, CA. 94583

A(! :+ +

EPA REG. NO. 201-357

CAUTION:

ACTIVE ING 2,2-dichic Related C INERT INGR *Equivalent



FOR LONG-LASTING FLEA CONTROL

Do not allow children to play with this collar — see cautions on back panel.

GREDIENTS lorovinyi dimethyi phosphate		BY WEIGHT
Compounds		0. 35°° *
REDIENTS		95.30°°
it to 4.7% w VAPONA: Insecticide.	TOTAL	100.00°5

CONTENTS: 1 FLEA COLLAR

NET WEIGHT: 0.4 OZ.

CLEAR