




CATTLE MINERAL BLOCK

(Contains RABON® Oral Larvicide)

NOT FOR HUMAN USE



TO PREVENT THE DEVELOPMENT OF HORN FLIES, FACE FLIES, HOUSE FLIES AND STABLE FLIES IN THE MANURE OF TREATED CATTLE.

ACTIVE INGREDIENTS	BY WEIGHT
2-chloro-1-(2, 4, 5-trichlorophenyl) vinyl dimethyl phosphate	1.8%*
INERT INGREDIENTS	98.2%**
TOTAL 100.0%	

GUARANTEED ANALYSIS	
CALCIUM (Ca)	Minimum 14.00%
CALCIUM (Ca)	Maximum 16.00%
PHOSPHORUS (P)	Minimum 4.50%
SALT	Minimum 30.00%
SALT	Maximum 34.00%

* Each pound contains 8.2 g RABON®
 ** Refers only to ingredients which are not larvicidal.
 EPA Reg. No. 201-393

INGREDIENTS
 Salt, Limestone, Steamed Bone Meal, Dicalcium Phosphate, Dried Molasses, Condensed Extracted Glutamic Acid Fermentation Product, Cane Molasses, Anise Oil, Fenugreek, Sulfur, Manganese Sulfate, Zinc Oxide, Calcium Iodate, Cobalt Carbonate.

CAUTION

KEEP OUT OF REACH OF CHILDREN

FEEDING DIRECTIONS

Provide one block per 15 – 20 head of cattle. Add an additional block when each block has been one-half consumed. Cattle should consume an average of 1/4 lb. of the block per 100 pounds of body weight per month. This will supply the recommended average intake of 70 mg RABON® Oral Larvicide per hundred pounds of body weight. Animals should not be starved for minerals before initiation of the RABON cattle Mineral Block program but OMIT all salt from other feed after the program is started. Do not feed free choice sources of salt or mineral containing salt in any form. Place near watering and loafing areas. Protect from rain.

If consumption of Shell Cattle Mineral Block is above 1/4 lb. of the block per 100 lbs. of body weight per month, reduce the number of feeding locations or relocate. If consumption is below 1/4 lb. of the block per 100 lbs. of body weight per month, increase the number of feeding locations. Start feeding Shell Cattle Mineral Block early in the spring before flies begin to appear, and continue feeding throughout the summer and into the fall.

Shell Cattle Mineral Block may be fed up to slaughter and to lactating dairy cows.

Shell Cattle Mineral Block prevents the development of horn flies, face

flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Shell Cattle Mineral Block should be used in conjunction with other good management and sanitation practices.

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse container. Destroy when empty.

40 POUNDS NET WEIGHT

SHELL CHEMICAL COMPANY, A DIVISION OF SHELL OIL COMPANY, ANIMAL HEALTH, SAN RAMON, CALIFORNIA