



# Pydrin<sup>®</sup>

## Insecticide

### 2.4 Emulsible Concentrate

**RESTRICTED USE  
PESTICIDE**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABEL**

PYDRIN<sup>®</sup> Insecticide 2.4 Emulsible Concentrate  
EPA Reg. No. 201-401

**DIRECTIONS FOR USE  
ON CHRISTMAS TREES, PINE SEED ORCHARDS AND  
FOREST TREE NURSERIES.**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Site	Insect	Usage			Further Use Directions	
		15 oz/100 gals	High Volume Sprays 11 oz/100 gals	1 1/2 - 2 1/2 gal of 2.4 EC per acre		Low Vol. Sprays (Mistblower) 1/2 oz/A
Christmas Tree Plantings, Pine Seed Orchards, Forest Tree Nurseries	Needle-bark Pine Tip Moth, Spittlebugs, Pineleaf Cuckoo, Redheaded Pine Sawfly, Balsam Woolly Agedid, Pine Chaffer, Red Pine Sawfly, Pine Conifer Bug	0.1 - 0.2	5 1/3 - 10 2/3	1 1/2 - 2 1/2 Teaspoon	0.1 - 0.2	Apply as needed to maintain control. Spray sufficient gallonage to obtain good coverage of entire tree. High volume sprays on Virginia Pine may cause some yellowing of needles. Low volume (Mistblower) applications at the rate of 0.1 to 0.2 lb a/acre are recommended to avoid this problem.
	Palm Weevil, Hortensia Leaf Weevil	0.1 - 0.2	5 1/3 - 10 2/3	1 1/2 - 2 1/2 Teaspoon		Apply as needed to maintain control. Dilute formulation in herbicide, instead of water, and spray tree stumps to control developing larvae. Use sufficient spray volume to thoroughly cover stump.
	Conoworm, Seedbug	11 oz/100 gals water for high volume sprays 3.5 pts/100 gals water for low volume sprays 0.25 lb a/acre apply, do not less than 10 gals of water for aerial applications				Apply first spray within 1 week of female flower closure or peak pollen flight for webbing Conoworm control. For other Conoworms and Seedbugs, apply first spray within 30 days following female flower closure. Repeat apply at intervals of 4 weeks but do not exceed 6.3 lb a/acre/yr.  Apply sprays: 5-10 gals of the 11 oz/100 gal dilution per acre with high volume sprayers; with low volume sprayers apply 100 gals of the 3.5 pts/100 gal dilution per acre.  Do not graze or harvest cover crop. Avoid contact with open water.

**ACCEPTED**

SEP 27 1984

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended for the pesticide  
registered under  
EPA Reg. No. 201-401