


1/22

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number 220135
	Application for Pesticide - Section I		

1. Company/Product Number 211-36	2. EPA Product Manager	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Tri-Cen	PM# 32	
5. Name and Address of Applicant (Include ZIP Code) Central Solutions, Inc. P.O. Box 15276 3130 Brinkerhoff Road Kansas City, KS 66115 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated <u>07 24 1996</u> <input checked="" type="checkbox"/> Notification - Explain below.	<div style="font-size: 1.5em; font-weight: bold; margin-bottom: 10px;">NOTIFICATION</div> <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below.
--	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of Addition of Indoor, Nonfood Sites for Antimicrobial Product. This notification is consistent w/ the provision of PR Notice 95-2 & EPA regs at 40 CFR 152.46 & no other changes have been made to the labeling or the CSF of the product. I understand that it's a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I further understand that if this notification isn't consistent w/ PR 95-2 & 40 CFR 152.46, this product may be in violation of FIFRA & I may be subject to enforcement action & penalties under sec **Section - III** 12 & 14 of FIFRA.

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container

3. Location of Net Contents Information
 Label Container

4. Size(s) Retail Container 1, 2, 4, 8, 16, 32, 64 oz; 5, 6, 15, 30, 55 gal. bulk


5. Location of Label Directions on label

6. Manner in Which Label is Affixed to Product.
 Lithograph Paper glued Stenciled Other silk-screened; sleeve labels

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name <u>Paul Nobrega</u>	Title <u>Director, Regulatory Affairs</u>	Telephone No. (Include Area Code) (913) 621-6542
--------------------------	---	---

<p style="text-align: center;">Certification</p> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Data Application Received (Stamped) SSG 1/8/97
2. Signature 	3. Title <u>Director of Regulatory Affairs</u>	
4. Typed Name Paul Nobrega	5. Date 10-04-96	

2/20

(CENTER PANEL)
(REQUIRED)

TRI-CEN

NOTIFICATION

301 2 4 1995

**GERMICIDAL DETERGENT
WITH ORGANIC SOIL TOLERANCE**

**BACTERICIDAL - TUBERCULOCIDAL - PSEUDOMONACIDAL
GERMICIDAL - FUNGICIDAL - VIRUCIDAL***

A highly effective combination odor counteractant,
soapless cleaner, and activated phenolic germicide

Formulated for effective health care facility sanitation
Formulated for effective hospital and health care facility sanitation
Formulated for health care facilities
Formulated for hospital and other health care facilities

Active Ingredients

Sodium ortho-benzyl-para-chlorophenate.....	4.40%
Sodium ortho-phenylphenate.....	2.82%
Sodium para-tertiary-amylphenate.....	2.49%
Inert Ingredients.....	90.29%

SEE BACK OR LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

KEEP OUT OF REACH OF CHILDREN

DANGER



CENTRAL SOLUTIONS, INC.

3130 BRINKERHOFF RD.
P.O. BOX 15276
KANSAS CITY, KANSAS 66115

EPA REG. NO. 211-36
EPA EST. NO. 211-KS-1

NET CONTENTS GAL.

3/22

(BACK OR LEFT PANEL)
(REQUIRED)

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE - Store in securely closed original container. Store in area that will prevent cross contamination of other pesticides, fertilizer, food and feed. Avoid storage at temperature extremes or in sunlight.

PESTICIDE DISPOSAL - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticides is in violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL - Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Reg. No. 211-36

EPA Est. No. 211-KS-1

ALTERNATE (Use when container size less than 1 gallon)

STORAGE AND DISPOSAL - Do not contaminate water, food or feed by storage or disposal.

STORAGE - Store in securely closed original container. Avoid storage at temperature extremes or sunlight.

PESTICIDE DISPOSAL - Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL - Do not reuse container. Replace cap and discard container in trash.

4/22

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CLEANING AND DISINFECTION

For broad spectrum disinfection of gram negative and gram positive bacteria, including *Pseudomonas aeruginosa*, add 1 ounce TRI-CEN to the gallon of water. Remove heavy soil or gross filth then apply solution with a mop, cloth, sponge, or hand pump trigger sprayer, so as to wet the surface thoroughly. Allow to remain wet for 10 minutes and then let air dry. Allow to remain wet for 10 minutes and then let air dry. Where Tuberculocidal activity is required, clean the surface prior to application of the disinfectant solution. If higher detergency is desired, increase dilution to 2 ounces per gallon of water. Prepare a fresh solution for each use.

(OR)

For broad spectrum disinfection of gram negative and gram positive bacteria, including *Pseudomonas aeruginosa*, add 1 ounce TRI-CEN to the gallon or water. Remove heavy soil or gross filth then apply solution with a mop, cloth, sponge, or hand-pump trigger sprayer, so as to wet the surface thoroughly. Allow to remain wet for 10 minutes and then let air dry. Where Tuberculocidal activity is required, allow the disinfectant solution to remain on the surface for 15 minutes. If higher detergency is desired, increase dilution to 2 ounces per gallon of water. Prepare a fresh solution for each use.

(Health Care Use)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

CLEANING AND DISINFECTION

For broad spectrum disinfection of gram negative and gram positive bacteria, including *Pseudomonas aeruginosa*, add 1 ounce TRI-CEN to the gallon of water. Remove heavy soil or gross filth then apply solution with a mop, cloth, sponge, or hand pump trigger sprayer so as to wet the surface thoroughly. Allow to remain wet for 10 minutes and then let air dry. Where Tuberculocidal activity is required, clean the surface prior to application of the disinfectant solution. If higher detergency is desired, increased dilution to 2 ounces per gallon of water. Prepare a fresh solution for each use.

(OR)

For broad spectrum disinfection of gram negative and gram positive bacteria, including *Pseudomonas aeruginosa*, add 1 ounce TRI-CEN to the gallon of water. Remove heavy soil or gross filth then apply solution with a mop, cloth, sponge, or hand pump trigger sprayer, so as to wet the surface thoroughly. Allow to remain wet for 10 minutes and then let air dry. Where Tuberculocidal activity is required, allow the disinfectant solution to remain on the surface for 15 minutes. If higher detergency is desired, increase dilution to 2 ounces per gallon of water. Prepare a fresh solution for each use.

EFFICACY PERFORMANCE

The broad spectrum efficacy of TRI-CEN has been confirmed using A.O.A.C. Protocol testing at a use-dilution of 1:128 in the presence of 5% blood serum (used to simulate organic soil load) at 20 degrees Celsius with an exposure time of 10 minutes.

GERMICIDAL:

TRI-CEN is effective against the broad spectrum of Gram positive and Gram negative pathogens as confirmed against *Salmonella choleraesuis*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa* by the AOAC Use Dilution test, in the presence of 5% organic soil load at 20 degrees Celsius with an exposure time of 10 minutes.

TRI-CEN is also effective against *Salmonella choleraesuis*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa* 28 days after dilution at 1:128 in the presence of 5% organic soil load at 20 degrees Celsius with an exposure time of 10 minutes following A.O.A.C. Protocol testing.

FUNGICIDAL:

At 1 ounce per gallon of water, TRI-CEN is also effective against *Trichophyton mentagrophytes*, (the athlete's foot fungus) in use areas such as shower room floors, locker room benches, bathmats or any other hard nonporous, inanimate environmental surfaces.

TUBERCULOCIDAL:

TRI-CEN disinfectant's effectiveness against *Mycobacterium bovis* (BCG) is demonstrated through testing using the AOAC Method "Confirmative in Vitro Test for Determining Tuberculocidal Activity" in 15 minutes at 20° C. Where Tuberculocidal activity is required, allow the disinfectant solution to remain on the surface for 15 minutes.

(Health Care Use)

VIRUCIDAL

***KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type-1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION:

Disposable gloves should be worn when cleaning blood, body fluids or contaminated items. Hands should be washed immediately and thoroughly after gloves are removed, or if hands are contaminated with blood or body fluids. Masks and protective eyewear or faceshields should be worn during cleaning procedures involving blood or body fluids. Gowns or aprons should be worn during cleaning procedures for protection against splashes of blood or other body fluids.

CLEANING PROCEDURE:

Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant. After cleaning is complete, follow disinfection directions described previously.

CONTACT TIME:

Leave surfaces wet for 10 minutes. Let air dry.

DISPOSAL OF INFECTIOUS MATERIALS:

Contaminated waste and blood and other body fluids should be autoclaved and disposed according to local regulations for infectious waste disposal.

(HAIR SALON USE)

TRI-CEN is recommended for use on brushes, combs, rollers, manicure, and other salon tools commonly found in beauty salons, barber shops, and hairstyling establishments.

TRI-CEN is highly effective for immersion disinfection of inanimate, non-porous surfaces. Use at a use dilution of 1:128 ($\frac{1}{4}$ ounce per 32 ounces of water, or 8 ml per liter).

DISINFECTION: Thoroughly clean and rinse instruments before immersion in disinfection solution. Immerse instruments completely for 10 minutes at room temperature (20°C/68°F). After immersion, remove instruments and rinse them thoroughly. Fresh solution should be prepared daily or more often if solution becomes visibly dirty.

CONFIRMATION: When tested with full immersion at room temperature for 10 minutes in a solution created as above, TRI-CEN was confirmed effective against *Mycobacterium tuberculosis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella choleraesuis* and *Trichophyton mentagrophytes*.

For container size of 16 fluid ounces:

This product makes 64 quarts of disinfectant solution when diluted $\frac{1}{4}$ oz. of disinfectant to 32 ounces of water (1 ounce of disinfectant to 128 ounces of water).

10/22

(MUSHROOM FARM PREMISE USE)

GERMICIDAL-DETERGENT WITH ORGANIC SOIL TOLERANCE
FORMULATED FOR MUSHROOM FARM SANITATION

TRI-CEN effectively
Kills fungi and bacteria which prove dangerous to mushroom crops.

DO NOT APPLY TO MUSHROOM CROP. DO NOT APPLY TO COMPOST OR CASING.
USE ONLY AS DIRECTED

EFFICACY PERFORMANCE

GERMICIDAL:

TRI-CEN has been tested and found effective against bacteria detrimental to the mushroom crop including:

- Pseudomonas fluorescens (Biotype G)....Bacterial Blotch Disease
- Pseudomonas aeruginosa.....Mummy Disease

FUNGICIDAL:

Fungi specific to the mushroom industry which TRI-CEN has been tested against and found effective include:

- Verticillium fungicola.....(Verticillium disease, Brown spot or Fungus spot)
- Mycogone perniciosa....(Mycogone disease or Wet bubble)
- Tricoderma viride.....(Tricoderma spot, Tricoderma blotch, Tricoderma mildew, Green mold)
- Dactylium dendroides...(Hypomyces mildew, Cobweb, Soft decay)

MUSHROOM FARM INDUSTRY USE

General Information

TRI-CEN is a phosphate free, germicidal detergent effective in the presence or moderate amounts of organic soil on a variety of surfaces found at mushroom farms. TRI-CEN's tri-phenolic formula is designed for one-step cleaning and disinfecting of hard inanimate non-porous environmental surfaces. TRI-CEN is strong enough to have no harmful effects on the surfaces being treated when used as directed.

Areas of Use

TRI-CEN has been designed for use between crops on surfaces that do not come in contact with the crop. Areas of intended use include tools, breezeways and track alleys before spawning, inside and outside walls of mushroom houses, lofts, floors, storage sheds, spawning machines, tampers and casing rings. Use of TRI-CEN should be limited to areas where compost and mushrooms are not present.

(POULTRY AND SWINE PREMISE USE)

Formulated for Effective
Poultry & Swine Premise Sanitation

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

SITE PREPARATION

The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated, disinfection can be accomplished in a single step.

CLEANING AND DISINFECTION

For all general cleaning and disinfection, use 1 ounce of TRI-CEN per gallon of water (1:128 solution). Apply TRI-CEN using a cloth, mop, or pressure sprayer so as to thoroughly wet surface to be disinfected. Allow to remain wet for 10 minutes and then let air dry. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use.

EFFICACY PERFORMANCE

The broad spectrum efficacy of TRI-CEN has been confirmed using A O A C protocol testing at a use dilution of 1:128 in the presence of 5% blood serum (used to simulate organic soil load) at 20 degrees Celsius with an exposure time of 10 minutes.

GERMICIDAL

TRI-CEN is effective against the broad spectrum of Gram positive and Gram negative pathogens as confirmed against *Salmonella choleraesuis*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa* by the A O A C Use Dilution Confirmation Test.

FUNGICIDAL

TRI-CEN's fungicidal efficacy has been confirmed through testing using the A O A C Fungicidal Test against the pathogenic fungi: *Trichophyton mentagrophytes*.

***VIRUCIDAL**

Virucidal testing using prescribed E P A testing procedures has verified TRI-CEN's effectiveness against the following viruses commonly associated with poultry and swine areas including:

Avian Influenza Virus	Avian Herpes Virus 2 (Marek's Disease)
Avian Bronchitis Virus	Hog Cholera Virus
Avian Laryngotracheitis Virus	Pseudorabies Virus

(POULTRY AND SWINE PREMISE USE)

AREAS OF USE
POULTRY PREMISE SANITATION (HATCHERIES)

- | | |
|---------------------|-----------------------------|
| -Egg Receiving Area | -Tray Dumping Area |
| -Egg Holding Area | -Chick Procession Area |
| -Setter Room | -Chick Holding Room or Area |
| -Hatchery Room | -Chick Loading Area |
| -Poultry Buildings | -Hauling Equipment |

APPLICATION AND USE DILUTIONS

POULTRY HOUSES:

1. Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water.
2. Use 1 ounce of TRI-CEN per gallon of water. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

HATCHERIES:

1. Use 1 ounce of TRI-CEN per gallon of water to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

EQUIPMENT:

1. Empty all troughs, rack, feeding and watering bins, buckets and fountains.
2. Use 1 ounce of TRI-CEN per gallon of water to treat all equipment. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

VEHICLES:

1. Clean all vehicles including mats, crates, cabs and wheels with high pressure water and TRI-CEN disinfectant. Use 1 ounce per gallon of water to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

SHOEBATHS:

1. Shoebath containing one inch of freshly made disinfectant solution should be placed at all entrances to buildings and hatcheries. Change the disinfectant solution in the bath daily or when solution appears dirty.

(POULTRY AND SWINE PREMISE USE)

POULTRY AND ANIMAL INDUSTRY USE

FOR CLEANING AND DISINFECTION OF

EQUIPMENT including feeders waterers fans racks troughs, bins and buckets VEHICLES including live haul poultry delivery truck feed trucks service person cars tractors and all other vehicles HATCHERIES including floors, walls, equipment, incubators hatchers, tray and buggies SHOEBATHS including all entrances to buildings and hatcheries

AREAS OF USE

SWINE AND CATTLE SANITATION

- | | |
|----------------------------|--------------------|
| -Farrowing Barns and Areas | -Waterer & Feeder |
| -Blocks | -Nursery |
| Creep Area | -Hauling Equipment |

MISCELLANEOUS

- | | |
|----------------------|---------------------------|
| -Horse Stables | -Sales Barn Stalls |
| -Dressing Plants | -Chutes/Loading Equipment |
| Sick Pens | -Veterinary Hospitals |
| -Cat and Dog Kennels | |

General Disinfection Uses

Can be used in the routine disinfection of hog houses, farrowing houses, finishing houses growers feeders and other equipment horse stables and sick pens at race tracks or sale barns hospital pens, calf pens and equipment Disinfecting houses used for poultry turkeys, ducks and game birds and for equipment used in brooding growing and laying quarters Disinfecting dog and cat kennels as well laboratory animal facilities at veterinary hospitals research laboratories and medical institutions Spraying cattle pens chutes loading equipment trucks and animal manure pits for the control of odor and ammonia

Dressing Plant Use

Disinfect equipment utensils walls and floors in poultry and animal dressing plants Equipment and utensils must be thoroughly rinsed with potable water before operations are resumed Disinfect offal rooms exterior walls and loading platforms of dressing plants

(POULTRY, SWINE AND FARM USE)

APPLICATION AND USE DILUTIONS

FARM PREMISE:

DO NOT USE IN MILKING STALLS, MILKING PARLORS OR MILK HOUSES.

1. Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.
2. Use 1 ounce of TRI-CEN per gallon of water (1 part TRI-CEN to 128 parts of water). Saturate all surfaces with the recommended solution for a period of 10 minutes.
3. Use 25-50 gallons of disinfectant solution per 1000 square feet of floor space. Totally immerse equipment to be treated in disinfectant solution. Ventilate building and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or aired. Thoroughly scrub all treated feed racks, troughs, automatic feeder, fountains, and waterers with soap or detergent and rinse with potable water before reuse.

EQUIPMENT:

1. Empty all troughs, racks, feeding and watering bins, buckets and fountains.
2. Use 1 ounce of TRI-CEN per gallon of water (1 part TRI-CEN to 128 parts of water) to saturate all equipment. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

VEHICLES:

1. Clean all vehicles including mats, crates, cabs, and wheels with high pressure water and TRI-CEN solutions.
2. Use 1 ounce of TRI-CEN per gallon of water (1 part TRI-CEN to 128 parts of water) to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

HATCHERIES:

1. Treat all surfaces thoroughly with TRI-CEN solution by scrubbing with brush under high water pressure with emphasis on cracks and moulding where fungus and mold are usually located.
2. Use 1 ounce of TRI-CEN per gallon of water (1 part TRI-CEN to 12 parts of water) to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution for 10 minutes or more.

SHOEBATHS SANITIZER:

1. Shoebath containing one inch of freshly made disinfectant solution should be placed at all entrances to buildings and hatcheries. Change the disinfectant solution in the bath daily or when solution appears dirty.

17/22

(AQUACULTURE USE)

Germicidal Detergent Formulated For Disinfecting Tools Equipment,
and Surfaces in Greenhouses Micropropagation Laboratories
Aquaculture and Hydroponic Farms Facilities or Plants

WARNING!!!

MUST NOT COME IN CONTACT WITH EDIBLE CROPS

GENERAL INFORMATION

TRI-CEN disinfectant is a phosphate free germicidal detergent effective in the presence of moderate amounts of organic soil on a variety of surfaces found in greenhouses micropropagation laboratories aquaculture and hydroponic farms TRI-CEN's multisynthetic phenolic formula is designed for one-step cleaning and disinfecting hard inanimate nonporous surfaces TRI CEN is strong enough to kill a broad spectrum of plant pathogens and pests yet mild enough to have no harmful effects on the surfaces being treated when as directed

AREAS OF USE

TRI-CEN is designed to be used between crops, or between daily work procedures on surfaces, tools and equipment that do not come in contact with edible crops Areas of intended use include tools floors walls, bench tops potting areas, headhouses dry storage areas sheds barns, automated potting equipment, cultivating and planting equipment pruning and budding equipment pallets and pallet moving equipment **DO NOT APPLY TO CROP SOIL OR GROWING MEDIA IN WHICH A CROP GROWS OR WILL BE GROWN!** To assure compliance rinse treated surfaces and tools potable water before they contact the crop or the substrate in which the crop is or will be growing

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

SITE PREPARATION

The first step in any on-going sanitation program should be the removal of gross contamination and debris This may be accomplished using a shovel broom, or vacuum depending on the area to be disinfected The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter Once the heavy debris eliminated disinfection can be accomplished in a single step

CLEANING AND DISINFECTION

For all general cleaning and disinfection use 1 ounce of TRI CEN per gallon of water (1 128 solution) Apply TRI CEN using a cloth mop or pressure sprayer so as to thoroughly wet surface to be disinfectant Allow to remain wet for 10 minutes and then let air dry For heavily soiled areas a precleaning step is required Prepare a fresh solution for each use

19/22

(INSTRUMENT CLEANING AND DECONTAMINATION USE)

DISINFECTANT CLEANER - FOR ULTRASONIC USE
INSTRUMENT PRESOAK
DISINFECTANT ULTRASONIC CLEANER CONCENTRATE

An effective multi-phenolic germicide for cleaning metal and/or ceramic instruments at chairside and immersion decontamination. **MAKES (X) GALLONS** - The contents of this bottle dilute to make (X) gallons. (X) liters cleaning and decontamination solution. **NOT RECOMMENDED FOR PLASTIC AND RUBBER.**

- ◆ Ideal as a chairside pre-soak for instruments
- ◆ Ideal as a holding solution prior to ultrasonic cleaning
- ◆ Prevents drying of blood, serum and debris
- ◆ Promotes safer instrument handling
- ◆ Won't discolor instruments or surface
- ◆ Pleasantly scented
- ◆ This product makes (X) gallons (X will depend on the size of container)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE:

TRI-CEN instrument pre-soak is an effective germicide detergent for immersion cleaning and/or decontamination of inanimate, non-porous surfaces on instruments and appliances prior to terminal sterilization or high-level disinfection. Use at a dilution of 1:128 (¼ ounce of TRI-CEN per quart of water or 8 ml per liter).

DILUTION:

Remove outer cap and seal. While holding bottle by its neck, twist off the inner cap and discard. Replace outer cap loosely. Gently squeeze the lower chamber until the upper reservoir is filled to the desired level (¼ oz per quart of water, or 8 ml per liter of water). Pour the concentrate into the tray or ultrasonic tank and add water to the recommended level.

CHAIRSIDE PRE-SOAK:

Place soiled instrument in holding tray solution until ready to be cleaned. If instrument is required again for the same patient, remove from holding solution and rinse thoroughly under gently running tepid water. Dispose of used solution after each patient. Rinse items prior to ultrasonic cleaning.

(INSTRUMENT CLEANING AND/OR DECONTAMINATION USE)

DECONTAMINATION:

Thoroughly clean and rinse instruments to remove excess debris, preferably in an ultrasonic cleaner before immersing in freshly prepared cleaning and/or decontamination solution. Cleanse and rinse the lumens of any hollow instruments before filling with solution. Avoid entrapment of air bubbles. All instruments and devices must be rinsed and rough dried before immersion in the solution. Immerse instruments completely for 10 minutes at room temperature (20°C/68°F) to destroy the broad spectrum of Gram positive and Gram negative bacteria, pathogenic fungi and Mycobacterium tuberculosis. After immersion, remove instruments and rinse them thoroughly prior to terminal sterilization or high level disinfection. Prepare a fresh holding solution daily, or sooner, if solution is visibly dirty.

For cleaning and/or decontamination (prior to terminal sterilization or high level disinfection) of inanimate, non-porous surfaces of dental and laboratory instruments and appliances in ultrasound equipment use TRI-CEN at a dilution of 1:128 (1 oz TRI-CEN per gallon of water), (¼ ounce per quart of water). Fill tank or beaker with water to recommended level. Fill upper reservoir of dispenser bottle to correct level to attain use dilution, and add to tank or beaker. Add precleaned instruments to solution so surfaces remain wet for 10 minutes. Always cover tank or beaker when unit is operating. Always turn off machine when removing items. Operate the ultrasonic cleaner no more than 15 minutes to minimize vaporization. Do not allow solution to overheat. Prepare a fresh solution daily, or sooner, when solution is visibly dirty. Do not mix with other chemicals. Clean and rinse tank daily.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

21/22

(CLEAN ROOM USE)

[Statements to be added only to labels of containers destined for clean rooms]

Product and container sterilized and distributed by _____.

Sterile pharmaceutical clean room formula

21/22

(CLEAN ROOM USE)

[Statements to be added only to label of containers destined for clean rooms]

TRI-CEN is recommended for use by pharmaceutical and biotechnology industries; medical device manufacturers; hospitals; and health care institutions that are dedicated to controlling the hazards of cross contamination in their establishments. For use on hard, inanimate surfaces in aseptic filling and gowning rooms, general manufacturing areas, or on machinery, tables, counters, laminar flow benches, floors, walls, stainless steel, porcelain, glass, and chrome.

FUNGICIDAL: At 1 ounce per gallon of water, TRI-CEN is also effective against Trichophyton mentagrophytes in use areas as aseptic filling and gowning rooms, and general manufacturing areas or on any other hard, non-porous, inanimate environmental surfaces.

This product is not to be used as a terminal sterilant / high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.