

NET CONTENTS

ARCADIAN

30 GALLONS

LIME SULFUR SOLUTION

WARNING — Keep out of reach of children

For Additional WARNING Information See Other Panel

USDA Reg. No. 218-30

10.59 Pounds Per Gallon at 60° F.

Test Baume 31° at 60° F.

ACTIVE INGREDIENT:
Calcium Polysulfide 29%

INERT INGREDIENTS 71%



ALLIED CHEMICAL CORPORATION
DEPARTMENT AG
40 RECTOR STREET
NEW YORK, N.Y., U.S.A.

ACCEPTED

JUL 17 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 418-30

DIRECTIONS

NOVEMBER-DECEMBER SPRAY for control of Coryneum Blight (Shot-hole) on Peaches and Nectarines; Use 7 gallons of Lime Sulfur Solution in 100 gallons of water.

DORMANT SPRAY on Apples, Pears, Peaches, Plums and Prunes, Almonds, Cherries and Nectarines; for control of San Jose Scale, Brown Apricot Scale, Soft Brown Scale, Greedy Scale and certain other scale insects: Use 10 gallons of Lime Sulfur Solution in 100 gallons of water.

For control of Mealybug, Bud Mites, certain mite eggs, Italian Pear Scale and Oystershell Scale and eggs of Fruit Tree Leaf Roller (also for control of scale insects listed above): Use a combination of 1 to 4 gallons of a light medium flowable oil emulsion plus 3 to 5 gallons of Lime Sulfur Solution in 100 gallons of water.

PRE-BLOSSOM SPRAY (Swollen or Green Bud Stage) on Apples and Pears to control Scab: Use 5 gallons of Lime Sulfur Solution in 100 gallons of water.

CLUSTER SPRAY on Apples and Pears for control of Scab, Pear Leaf Blister Mite and certain insects such as the Bryobia Mite: Use 2 gallons of Lime Sulfur Solution plus 5 lbs. of Dritomic Sulfur (or Wettable Sulfur) in 100 gallons of water.

CALYX SPRAY on Apples and Pears to control Scab: Use 1 gallon of Lime Sulfur Solution in 100 gallons of water.

JACKET SPRAY on Peaches and Nectarines to control Powdery Mildew: Use 1/2 to 1 gallon of Lime Sulfur Solution plus 4 lbs. of Dritomic Sulfur (or Wettable Sulfur) in 100 gallons of water. For Silver Mite on Peaches and Nectarines: Use 2 gallons of Lime Sulfur Solution plus 4 lbs. of Dritomic Sulfur (or Wettable Sulfur) in 100 gallons of water.

POST-HARVEST SPRAY on Pears for control of Blister Mite: Use 2 gallons of Lime Sulfur Solution plus 4 to 5 lbs. of Dritomic Sulfur (or Wettable Sulfur) in 100 gallons of water.

NOTICE: Do not allow this material to freeze.

Do not apply Lime Sulfur sprays when freezing temperatures are predicted or when weather conditions may result in slow drying of sprays.

Do not apply Lime Sulfur sprays when temperature is over 90° F.

Do not apply oil sprays on apples within 3 weeks after applying Lime Sulfur sprays during growing season.

Do not use Lime Sulfur or Sulfur on Apricots.

WARNING! Contact with acid liberates poisonous gas. May burn skin or eyes.

Avoid contact with skin and particularly eyes.

Remove and wash contaminated clothing.

Keep away from acids.

In case of contact, immediately flush skin or eyes with plenty of water; for eyes get medical attention.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MADE IN U.S.A.

9441-1
1/305

PRINTED IN U.S.A.