

DIRECTIONS

ALL RATES ARE FOR USE IN 100 GALLONS OF WATER

MIXING AND APPLICATION: Mix the recommended amount of Dichlone in water until the spray mixture is uniform. Thoroughly wet plants with spray mixture.

APPLE: APPLE SCAB: Use ¼ pound in protectant sprays. Apply at 5 to 7 day intervals, from delayed dormant up to 10 days after petal fall. For eradicant sprays (up to 50 hours after infection period begins), use ½ pound. Do Not use Dichlone in sprays later than 1 day before harvest.

PEACH: BROWN ROT BLOSSOM BLIGHT: Use ½ pound. Apply sprays at pink, full bloom and just before shuck fall. Do Not apply later than 7 days before harvest.

LEAF CURL: Use 1 pound in a dormant spray just before buds begin to swell.

CORYNEUM BLIGHT (CALIFORNIA BLIGHT): Use ¾ pound between September 15 and September 30 in Oregon and Washington.

PLUM AND PRUNE: BROWN ROT BLOSSOM BLIGHT: Use ½ pound at pink, full bloom and pre-shuck fall stages of growth. When cool, wet weather occurs, resulting in a prolonged bloom period, apply Dichlone every 4 days.

BROWN ROT OF FRUIT: Use ½ pound at 7 to 10 day intervals beginning at shuck fall and continuing until just before harvest. Do Not apply to plums or prunes within 3 days of harvest.

CHERRY: BROWN ROT BLOSSOM BLIGHT: Use ½ pound. Apply at popcorn, full bloom and petal fall stages. Do Not apply later than 3 days before harvest.

TOMATO: LEAF BLIGHTS: Use ¾ pound. Apply sprays at 7 to 10 day intervals during infection periods.

PHOMA, GREY MOLD AND LATE BLIGHT (PLANT BED TREATMENT): Use 1 pound. Apply sprays at 5 to 7 day intervals during infection periods.

TRANSPLANTING TREATMENT: Use 1 pound per 100 gallons as a dip after pulling and before planting. Dip entire plant.

CELERY: BLIGHT: Use ½ pound. Apply 200 gallons of spray mixture per acre at 7 to 10 day intervals during infection periods.

ORNAMENTALS: BLACK SPOT OF ROSES: Use ¼ pound. Apply sprays when leaves first appear and repeat at 7 to 10 day intervals.

SMALL VOLUME RATES

½ level teaspoonful per gallon equals ¼ pound per 100 gallons.

1 level teaspoonful per gallon equals ½ pound per 100 gallons.

1½ level teaspoonfuls per gallon equals ¾ pound per 100 gallons.

2 level teaspoonfuls per gallon equals 1 pound per 100 gallons.

COMPATIBILITY: Dichlone is compatible with most common fungicides and wettable powder insecticides. Do Not use in sprays containing lime, calcium arsenate, mercury or dinitro compounds, oil, or emulsifiable concentrates.

NOTE: Do Not use more than the following amounts of this material per acre per application; on tomatoes 2½ pounds and on celery 1½ pounds.

CAUTION

MAY CAUSE SKIN IRRITATION!

Avoid prolonged or repeated contact with skin.

Avoid breathing dust or spray mist.

Use with adequate ventilation.

Wear protective clothing.

Wash with soap and water after using and before eating or smoking.

Avoid contamination of feed and foodstuffs.

NOTE: Skin irritation caused by Dichlone is similar to sunburn. It is most likely to occur in warm temperature conditions.

Toxic to fish. Keep out of any body of water.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use. Storage and handling of this material not in strict accordance with directions given herewith.

ARCADIAN®

NET WEIGHT 2½ POUNDS

DICHLONE

WETTABLE POWDER FUNGICIDE

CODE 9540

USDA REG. NO. 218-349

ACTIVE INGREDIENT:

Dichlone (2,3-dichloro-1,4-naphthoquinone) 50%

INERT INGREDIENTS

..... 50%



ALLIED CHEMICAL CORPORATION
DEPARTMENT AG, 40 RECTOR STREET, NEW YORK, N.Y., U.S.A.

CAUTION KEEP OUT OF REACH OF CHILDREN.
For Additional Caution Information See Back Panel.

PRINTED IN U.S.A.

9540
A1508E

MADE IN U.S.A.

