

ARCADIAN

NET WEIGHT 50 POUNDS

KILMAG®

A SPECIAL FORMULATION OF CALCIUM ARSENATE

For Control of Certain Fly Maggots
Under Poultry and Mink Cages,
and Turkeys on Wire Floors

CODE 9415

USDA Reg. No. 218-415

ACTIVE INGREDIENT: Calcium Arsenate 83.10%
INERT INGREDIENTS: 16.90%

TOTAL ARSENIC:
Expressed as Metallic Arsenic (As) 31.29%

ARSENIC IN WATER SOLUBLE FORM:
Expressed as Metallic Arsenic (As) 0.98%

POISON

ALLIED CHEMICAL CORPORATION
DEPARTMENT AG
40 RECTOR STREET
NEW YORK, N.Y., U.S.A.

NOTICE: This product is licensed for use only in accordance with the instructions on the label and not for the formulation of other pesticidal compositions.

WARNING!

POISONOUS IF SWALLOWED.

Do not breathe dust or spray mist.

Do not get on skin, in eyes or on clothing.

Wash thoroughly after using. Store away from feed and face cloths.

Completely empty container and dispose of pesticide by burying. Bury container immediately and stay out of smoke. Bury sales and residue.

POISON

DANGER

Keep out of reach of children and domestic animals.

CALL A PHYSICIAN IMMEDIATELY.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MADE IN U.S.A.

9415
A 1311A

PRINTED IN U.S.A.

FOR CONTROL OF THE LARVAL (maggot) STAGE OF THE HOUSE FLY AND SOLDIER FLY IN ACCUMULATIONS OF DROPPINGS UNDER CAGED BIRDS AND MINK AND TURKEYS ON WIRE FLOORS.

HOW TO MIX: Use 2 pounds in 4 to 6 gallons of water. When manure is wet use 4 gallons of water, when it is dry use 6 gallons. Mix well and agitate occasionally while spraying.

HOW TO APPLY: Apply the above amount evenly over 1000 square feet of manure as a coarse spray.

WHEN TO APPLY: Make a thorough overall application for first treatment. Check daily for one week and treat misses or areas where maggots are present. Then spray thoroughly every seven days to maintain control.

CAUTION: Do not apply manure treated with this product to food crops not covered by tolerances established under the provisions of the Federal Food, Drug and Cosmetic Act. Crops which have Established Tolerances for Calcium Arsenate and on which KILMAG-treated manure may be used are listed below:

- | | | |
|------------------|---------------|------------------------------|
| Asparagus | Collards | Pumpkins |
| Blackberries | Cucumbers | Raspberries |
| Black-eyed peas | Eggplant | Rutabagas |
| Blueberries | Huckleberries | Spinach |
| Broccoli | Kale | Squash (incl. summer squash) |
| Brussels sprouts | Kohlrabi | Strawberries |
| Cabbage | Loganberries | Tomatoes |
| Carrots | Melons | Turkeys |
| Caiflower | Peppers | |
| Celery | | |

Do not apply manure treated with this product to soils where rice, lima beans, snap beans or other crops which are sensitive to calcium arsenate are growing or may be planted later.

