

**WARNING!**

Hazardous if swallowed or absorbed through the skin.

Do Not get on skin, in eyes, or on clothing.

Wear rubber gloves when handling this product.

Wash with soap and water after using and before eating or smoking.

Wash contaminated clothing before re-use.

Do Not contaminate feed or foodstuffs.

Keep out of reach of children and domestic animals.

Do Not apply in areas accessible to children or domestic animals.

**DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY.**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**NOTICE:** This product is licensed for use only in accordance with the instructions on the label and not for the formulation of other pesticidal compositions.

MADE IN U. S. A.

9412-2  
A 1702 B

PRINTED IN U.S.A.

NET WEIGHT



25 POUNDS

CODE 9412

NEW IMPROVED FORMULATION . . .

**KEPONE®**

**ANT and ROACH BAIT • PELLETTED**  
FOR CONTROL OF CERTAIN ANTS AND ROACHES

**ACTIVE INGREDIENT:**

Decachlorooctahydro-1,3,4-metheno-2H-cyclobuta[cd]pentalen-2-one . . . . . 0.125%

**INERT INGREDIENTS** . . . . . 99.875%




**ALLIED CHEMICAL CORPORATION**  
AGRICULTURAL DIVISION  
MORRISTOWN, N. J., U.S.A.

**WARNING: Keep Out of Reach of Children**

For Additional WARNING Information See Other Panel

U.S. Pat. Reissue 24,749

**EPA Reg. No. 218-553**



**DIRECTIONS**

FOR USE IN RESTAURANTS, PLANTS, HOTELS AND

**ANTS**

For Harvester Ants, Black Cornfield Ant, Thief Ant, etc. Scatter Kepone bait within 6 feet of the colony may take 2 weeks. Treatment should be repeated. For Millipedes and Pillbugs, scatter bait on foundation of building.

**ROACHES AND SILVER**

For American, German, Brown Banded Roach, etc. Scatter 2 to 3 tablespoons of bait in suitable containers, out of reach of children or pets.


Kepone bait is best used as the only roach control. Do Not use Kepone bait in areas where it is used as the only roach control. Do Not use Kepone bait in areas where it is used as the only roach control.

**FOOD PROCES**

When plant food is not available, bait boxes can be accounted for to resuming

bed through  
 n clothing.  
 ng  
 using and  
 fore re-use.  
 dstuffs.  
 nd  
 to children or  
 WHEN EMPTY.  
 f application, weather and  
 specifically recommended,  
 duct are beyond the control  
 or handling of this material  
 with.  
 ordance with the instructions  
 sticidal compositions.  
 PRINTED IN U.S.A.

NET WEIGHT



25 POUNDS

CODE 9412

NEW IMPROVED FORMULATION . . .

# KEPONE®

ANT and ROACH BAIT • PELLETTED  
 FOR CONTROL OF CERTAIN ANTS AND ROACHES

**ACTIVE INGREDIENT:**

Decachlorooctahydro-1,3,4-metheno-2H-  
 cyclobuta[cd]pentalen-2-one . . . . . 0.125%

**INERT INGREDIENTS** . . . . . 99.875%




ALLIED CHEMICAL CORPORATION  
 AGRICULTURAL DIVISION  
 MORRISTOWN, N. J., U.S.A.

**WARNING: Keep Out of Reach of Children**

For Additional WARNING Information See Other Panel

U.S. Pat. Reissue 24,749

EPA Reg. No. 218-553



**DIRECTIONS**

FOR USE IN AND AROUND HOMES, RESTAURANTS, STORES, FOOD PROCESSING PLANTS, HOSPITALS, NURSING HOMES, HOTELS AND MOTELS.

**ANTS**

For Harvester Ants, Black Crazy Ant, Little Black Ant, Pyramid Ant, Pharaoh Ant, Cornfield Ant, Thief Ant, and Certain Other ANTS scatter 1 to 2 tablespoons of Kepone bait within 6 to 12 inches of the nest entrance. Additional bait may be scattered in the foraging area and around the foundations of buildings. Control of the colony may take 2 to 6 weeks. Repeat treatment after 6 weeks if necessary. Treatment should be made only when ants are active.

For Millipedes and Pillbugs scatter the bait in the foraging area and around the foundation of buildings. Repeat treatment as needed.

**ROACHES AND SILVERFISH**

For American, German, Oriental, and Brown-headed Roaches and for Silverfish use 2 to 3 tablespoons of Kepone bait for each infested room. Scatter bait, or place in suitable containers, behind or under fixtures and in other areas not accessible to children or pets.

Kepone bait is best employed as a supplemental residual treatment preceded by recommended roach sprays and good sanitation practices. When Kepone bait is used as the only roach treatment, control may not be obtained for 3 days to 4 weeks, depending upon conditions. Re-treat as necessary to effect and maintain control. Do Not use on or around food or forage crops.

**FOOD PROCESSING PLANTS:** Use only when plant is not in operation and food is not exposed. Use only in bait boxes where bait in each box can be accounted for. Remove and account for all bait boxes prior to resuming plant operation.