

NON-CROP LAND

For use on non-crop land such as Right-of-Ways, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas and Parking Lots for control of most annual and perennial grasses and broad-leaved weeds. Control will usually last 8 months or longer. The effectiveness and length of time control will last depend on type of soil, amount of rain and other conditions.

RATE: Under ordinary conditions use at the rate of 400 to 800 pounds per acre. Under conditions where weeds are difficult to control increase the rate to 1200 pounds per acre. For spot treatment use 1 to 2 pounds per 100 square feet. For uniform control spread the material evenly over the area to be treated.

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. Do Not apply when ground is frozen.

FLORIDA CITRUS GROVES

For Use in <u>Drainage Ditches</u> For Control of Most Annual and Perennial Grasses² and Broad-Leaved Weeds such as Paragrass, Arrowhead, Guineagrass and Water Primrose.


RATE: Use at the rate of 400 to 800 pounds per acre. See chart:

Amount to Aprily Per 1/2 Linear Feet of Ditch

Width of Ditch feet 2 4 6 8 10

Pounds Urox 4.5 2 to 4 4 to 5 6 to 12 8 to 16 10 to 20

(Continued on right panel)



NET WEIGHT 50 POUNDS

UROX® 5.5

WEED KILLER

For General Weed Control on Non-Crop Lands

CODE 9947

USDA Reg No. 218-631

ACTIVE INGREDIENT:

Monuron Trichloroacetate 13-(p-chlorophenyl)-1, 1-dimethylurea trichloroacetate 5.5°.

INERT INGREDIENTS:

11 S. Patent No. 2,782 112

94.5%

DEPARTMENT AG 40 RECTOR STREET NEW YORK, N.Y., U.S.A.

ALLIED CHEMICAL CORPORATION

- Keep out of reach of children

For Additional CAUTION Information See Right Panel

Made n USA



.

(Continued from left panel)

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. Treat only bottom not banks.

when to apply: Apply when ditch is not in use but soil is moist. If ditch bottom is not moist, wet it down before treating. If less than 4 inches of rain falls between time of treatment and time of intended use, run water into the ditch and let stand for 3 days. Drain off and waste the water before using ditch.

NOTE: Do Not use in ditches into which tree or other desirable plant roots extend. Do Not contaminate domestic water supplies, farm ponds or water to be used for livestock.

HARMFUL IF SWALLOWED

IRRITATING TO SKIN, EYES, NOSE AND THROAT

Do not breathe dust.

Do not get on skin, in eyes or on clothing. In case of contact, immediately wash skin with plenty of water; for eyes, wash with water for at least 15 minutes and get medical attention.

NOTE: Do not allow to drift onto desirable plants or cropland.

Do not apply rear lawns, valuable trees or other desirable plants or to areas where their roots may extend. Equipment used for application should be thoroughly cleaned to prevent corrosion or contamination of other materials used in it later.

Fortow directions carefully. Timing and method of application, weather and crop conditions mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Super assumes all risks of use storage or handling of this maternal not in storal accordance with directions given herewith

