



contains

ROTENONE 5%

For use in sprays to control certain

GARDEN INSECTS

ACTIVE INGREDIENTS-

Rotenone	and a community of the	5.0%
Other Cube Resins		7.5 ℃
INERT INGREDIENTS	يوري	8 7.5 %

CAUTION: Keep Out of Reach of Children See other Cautions on back panel

EPA Reg. No. 226-17

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin. Wear clean clothing. Avoid inhalation of this dust, as it may cause a temporary numbness of the respiratory tract. Breathing fresh air relieves this condition.

Keep out of any body of water. This product is poisonous to fish.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS

GARDEN SPRAY for control of Mexican Bean Beetle on Beans; Imported Cabbage worms and diamond back Moths on Cabbage; also pea aphids on Peas: Use 1 lb. to 25 gals, of water (6 level tablezpoonsful to 2 gals.). For home garden use spray plants thoroughly to upper and lower leaf surfaces when insects first appear. Repeat in 7 to 10 days or as often as necessary. When using on cabbage add 1 oz. soap powder for each gallon of water used. Do not apply to food crops later than 1 day of harvest.

For beans use up to 20 lb. per acre. For cabbage use up to 10 lb. per acre. For peas use up to 12 lb. per acre.

For a minimum dosage on beans, cabbages, and peas use 5 lb. per acre.

Tobacco States Chemical Co., Inc. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

#4 12571