

NET WEIGHT ONE POUND

Tobacco States

# 75% ZINEB WETTABLE POWDER

FOR CONTROL OF CERTAIN DISEASES  
ON TOBACCO, TOMATOES,   
 FLOWERS AND CERTAIN OTHER  
PLANTS.

**ACTIVE INGREDIENT:**

Zineb (zinc ethylene  
bisdithiocarbamate) . . . 75%

INERT INGREDIENTS . . . . . 25%  
100%

Caution: May cause irritation of nose, throat, and skin. Do not breathe dust or spray mist. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure. Keep container closed. Store in a dry place. Keep away from fire and sparks.

Do not reuse empty containers. Destroy them by burying with waste or burning. Stay away from smoke or fumes.

**CAUTION:** Keep Out of Reach of Children  
See other Cautions on back panel

**DIRECTIONS FOR USE**

**GENERAL DIRECTIONS**—Use 1½ to 2 pounds per 100 gallons of water (unless otherwise specified) in conventional sprayers, or equivalent in concentrate or aerial sprayers. Three pounds should be applied per acre on average size field crops.

Apply at first sign of, or prior to appearance of disease. Repeat at one or two week intervals as necessary unless otherwise noted.

**BEANS**—Downy Mildew, Rust, Anthracnose: Begin when first blossoms appear. Repeat applications at 4 to 7 day intervals. Do not apply within 7 days of harvest.

**CABBAGE, BROCCOLI, CAULIFLOWER, COLLARDS**—Downy mildew, Alternaria leaf blight: Do not apply within 7 days of harvest.

**CANTALOUPE, CUCUMBERS, WATERMELONS** and other **CUCURBITS**—Downy mildew, Alternaria (Macrosporium) blight, Anthracnose, Gummy stem blight: Direct spray especially at under-side of leaves, begin applications when plants appear above the ground.

**TOMATOES**—Early and late blight, Anthracnose, Stemphylium and Septoria leaf spots, Cladosporium gray mold: Begin spraying when disease is reported in area, and repeat applications at 7 to 10-day intervals or oftener if blight conditions are severe.

**STRAWBERRIES**—Leaf spot: Begin when new growth starts in spring. Repeat until fruit begins to form. Leaf scorch: Apply in new beds in late summer—2 to 3 weekly applications. Do not apply within 7 days of Harvest.

**TOBACCO**—Blue mold: Begin applications when plants emerge or when blue mold is first reported in area. Spray regularly 2 or 3 times per week during active blue mold weather. Repeat applications immediately after a rain. Use 3 lbs. 75% Zineb per 100 gal. water (2 level tablespoons per gallon). Apply at 3 to 6 gal. per 100 sq. yds. of plant bed, depending on size of plants.

**SNAPDRAGONS, CHRYSANTHEMUMS, CARNATIONS, HOLLYHOCKS**—Rust, Leaf spot and Anthracnose: Apply at weekly to 10 day intervals as soon as disease conditions threaten. For control of Botrytis petal spot and Ascochyta ray make application twice weekly during the blooming season.

**ROSES**—Black spot: Apply at weekly intervals. Applications should begin when new leaves unfold in spring

USDA Reg. No. 225-21

H 6-70

ACCEPTED  
UNDER THE FEDERAL INSECTICIDE ACT  
FUNGICIDE AND MITICIDE ACT  
FOR THE CONTROL OF FUNGUS AND MITES  
EDUCATIONAL INFORMATION  
TO THE TRADE COMMUNITY

PESTICIDE

TOBACCO STATES CHEMICAL CO., INC.  
LEXINGTON, KENTUCKY