

[226-3]

NET CONTENTS-1 POUND



## 1.00 % ROTENONE GARDEN DUST

for the control of

**MEXICAN BEAN BEETLES**

AND CERTAIN OTHER GARDEN INSECTS

### ACTIVE INGREDIENTS—

Rotenone . . . . . 1.00%

Other Cube Resins . . . . . 1.75%

INERT INGREDIENTS . . . . . 97.25%

**CAUTION:** Keep out of the reach of children.  
See other cautions on rear panel.

TOBACCO STATES CHEMICAL CO  
LEXINGTON KENTUCKY

# Tobacco States Pesticides

### CAUTION

Avoid prolonged inhalation of this dust as it may cause a temporary numbness of the respiratory tract. Fresh air relieves this condition.

Keep out of reach of young children. This product is dangerous to fish.

### DIRECTIONS

**NOTE:** Do not apply to food crops within one day of harvest.

**BEANS:** The Mexican Bean Beetle: This insect feeds from under side of leaves. Use duster with adjustable nozzle that will direct dust upward, from underneath plant. Use up to 100 lbs. per acre.

**CARROTS, BROCCOLI:** Cabbage worms, Diamond back Moth. Use up to 50 lbs. per acre.

**PEAS:** Pea Aphids. Use up to 60 lbs. per acre.

Better control can be obtained if plants are covered thoroughly by dusting at the first appearance of insects. Since this dust loses its toxicity in 7 to 10 days after being exposed to sunlight, dustings should be repeated at least every 10 days or more if necessary. Dusting should be done when foliage is moist with dew.

### FOR SPRAYING:

Add 1 lb. of this dust with about 1 ounce of soap powder to 3 1/2 gals. water. Mix thoroughly and spray plants every 10 days for good control.

6-3-67

USDA Reg. No. 226-31

ACCEPTED  
July 17, 1968  
226-31  
Pesticides  
Tobacco States Chemical Co.  
Lexington, Kentucky