

NET CONTENTS—2 POUNDS



TOMATO

COPPER

DUST

2000

for use as a dust
in control of

EARLY AND LATE BLIGHT ON
TOMATOES AND POTATOES

ACTIVE INGREDIENTS:

Copper, expressed as metallic 7.0%
(in Basic Copper Sulphate)

INERT INGREDIENTS 93.0%

CAUTION: Keep out of the reach of children.

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON KENTUCKY

DIRECTIONS

HOW TO APPLY: Apply TOBACCO STATES TOMATO COPPER DUST as it comes from the package with a hand duster. Apply lightly and evenly to both sides of leaves and stems. It is not necessary to discolor the foliage with dust; a light covering evenly applied is adequate for control. Apply more often in rainy weather. It is necessary to have the dust on the plants before diseases appear.

TOMATO—Controls Late Blight, Early Blight, Septoria Leaf Spot, Anthracnose, Leaf Mold. Start dusting when the first true leaves appear and repeat every 7 to 10 days. Apply within 24 hours after washing rains.

POTATOES—Controls Early Blight, Late Blight. Start dusting when plants are 6 inches high and repeat at 7 to 10 day intervals throughout the season.

CUCUMBERS, SQUASH, MELONS, CANTALOUPE—Controls Anthracnose, Downy Mildew, and Angular Leaf Spot. Dust lightly as soon as plants come through the ground and repeat at regular 7 to 10 day intervals. Do not make extremely heavy applications especially on young plants because extra heavy amounts of copper tend to injure some plants.

BEANS—Controls Downy Mildew, Anthracnose, and Angular Leaf Spot. Dust as soon as plants reach 5 inches height and continue weekly. For Mexican Bean Beetle control use TOBACCO STATES ROTENONE DUST.

For other uses and timing consult State Agricultural Experiment Station or State Extension Service Specialist.

TOBACCO STATES CHEMICAL CO., INC.

LEXINGTON, KENTUCKY

USDA Reg. No. 226-32