

ONE POUND



TRI - BASIC COPPER SULPHATE

for control of
EARLY AND LATE BLIGHT ON TOMATOES
AND POTATOES

for use as a spray to be
mixed with water

(COMMERCIAL 4:1 BASIC COPPER SULPHATE)

ACTIVE INGREDIENT(S)

Copper, expressed as metallic

43%

(in the form of Basic Copper Sulphate)


INERT INGREDIENT(S)

47%

WARNING: Keep Out of Reach of Children

See other Warning on back panel

HARMFUL IF SWALLOWED



WARNING: HARMFUL IF SWALLOWED.

DIRECTIONS FOR TOBACCO PLANT BEDS (Ky., Tenn., Ohio, Indiana):
Wildfire and Angular Leafspot: Mix 1/2 lb. Tri-Basic Copper with 25 gallons of
water for treatment of 100 sq. yds. (9 x 100 ft. or 12 x 75 ft.) Apply with a sprink-
ling can. 1st application when plants are up (two-leaf stage). 2nd application again
in 7 to 10 days. (Tri-Basic Copper is ready to use without lime.)

DIRECTIONS FOR SPRAYING TENNESSEE TRI-BASIC

CROP	TO CONTROL	TRI-BASIC PER 100 GALLONS OF SPRAY
TOMATO	Late Blight, Early Blight, Septoria Leaf Spot, Leaf Mold and Anthracnose	4 pounds - Apply both in seed beds and field
POTATO	EARLY AND LATE BLIGHT	4 pounds
CUCUMBERS, SQUASH, MELONS, CANTALOUPE	ANTHRACNOSE, Downy Mildew, Angular Leaf Spot	2 pounds
CABBAGE, BROCCOLI, CAULIFLOWER	Downy Mildew, Leaf Spot	3 pounds
BEANS	Downy Mildew, Anthrac- nose, Angular Leaf Spot	2 pounds - start when plants are about 5 inches high
GRAPES	Black Rot, Downy Mil- dew, Anthracnose	2 pounds

2 oz. (5 tablespoons) in 3 gallons is equivalent to 4 lbs. per 100 gals.

FOR STICKER: Add 1/2 lb. of flour to each 1 lb. of Tri-Basic or 1/4 oz. (4
tbls.) to each 2 oz. Tri-Basic.

For use as a dust use commercially prepared dust which have a better blend
such as TOBACCO STATES TOMATO COPPER DUST. For timing and other
uses, consult State Agricultural Experiment Station or State Extension
Service authorities.

TRI-BASIC should be mixed with water and applied every 7 to 10 days and
after washing rains.

USDA No. 226-33

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY