

NET WEIGHT ONE POUND

**Tobacco States**

# 10% B-H-C

**Wettable Powder**

**An Insecticide Containing 10%  
of the Gamma Isomer of  
Benzene Hexachloride**

ACTIVE INGREDIENTS  
Gamma Isomer of Benzene Hexachloride 10.0%  
Other Isomers of Benzene Hexachloride 1.0%  
INERT INGREDIENTS 89.0%

## WARNING

Keep out of reach of children.  
Harmful if swallowed. Harmful vapor and dust may cause irritation of the skin and eyes. May be absorbed through the skin. Avoid inhaling dust, vapor or mist from the spray. Avoid contact with the skin and eyes. In case of skin contact, wash with plenty of mild soap and water. For eyes, flush with water for 15 minutes. Get prompt medical attention.  
Do not use on dairy animals or beef cattle. Avoid contamination of milk and foodstuffs and feed.

EPA Reg. No. 226-38

**ACCEPTED**  
226-38  
Dec 14, 1970  
UNDER THE FEDERAL INSECTICIDE ACT  
FUNGICIDE, PESTICIDE AND POLYMER REGISTER  
EPA REG. NO. 226-38

## DIRECTIONS FOR USE

When using as a spray, no pre-slurrying is necessary. Merely stir the required amount of 10% BHC into the required amount of water.

ONE TEASPOONFUL OF 10% BHC IN 1 GALLON OF WATER IS EQUAL TO 1 LB. IN 100. NOTE: Consult the Agricultural Experiment Station or Extension Service Specialist regarding the number and frequency of applications. Do not use on poultry, dairy animals or feed to be fed to dairy animals being fed for 14 days after. Do not use in or on poultry houses. Do not use in dairy barns or milk rooms.

**APPLE, Aphids** — Use 2 lbs. 10% BHC in 100 gal. water. May be used as late as petal fall at the 2 lb. dosage for the control of Rosy Apple Aphids. Do not apply more than 36 lb. per acre application. Do not apply within 60 days of harvest or not after fruit starts to form.

**PEACH, Plum Curculio** — Use 2 lbs. 10% BHC in 100 gal. water. Do not apply more than 36 lb. per acre application. Make first application when 1/2 of shucks have fallen. Second application when all of shucks have fallen. Third application about 7 days later. Do not use within 60 days of harvest.

**STRAWBERRIES, Spittle Bugs, Strawberry Weevils** — Use 2 lbs. 10% BHC in 100 gal. of water. Do not apply more than 36 lb. per acre application. Do not apply after the edible parts start to form.

## OBSERVE THE FOLLOWING PRECAUTIONS

- Use only as directed. Do not apply to the water in any container.
- DO NOT APPLY to any surface where there is a possibility of contact with food.
- DO NOT USE on any surface where there is a possibility of contact with food.
- DO NOT APPLY to any surface where there is a possibility of contact with food.
- Do not apply to any surface where there is a possibility of contact with food.
- Do not apply to any surface where there is a possibility of contact with food.
- Apply on a calm day or evening with the wind at or near land.
- Toxic to Fish and Wildlife; do not contaminate ponds, lakes, or streams with this material.

16-70

**PESTICIDE**

**TOBACCO STATES CHEMICAL CO., INC.**  
LEXINGTON, KENTUCKY