

CONTENTS ONE GALLON

Tobacco States
CHEMICAL CO., INC.

TOXAPHENE

6 lb. Emulsion Concentrate

For dilution with water to form emulsions suitable for control of
GRASSHOPPERS, ARMYWORMS and GREEN CUTWORMS

ACTIVE INGREDIENTS

Toxaphene-Technical Chlorinated Camphene
(67%-69% Chlorine content) 60.2%

Petroleum distillates 35.8%

INERT INGREDIENT: 4.0%

(Contains 6 lbs. Technical Toxaphene per gal at 70°F)

WARNING: Keep out of reach of children.

MAY BE FATAL IF SWALLOWED. Do not breathe spray mist. Do not get on skin, in eyes, or on clothing. Wash thoroughly after use. Avoid contamination of feed and food. Not for use or storage in or around the home. See other required warning statements on the back label.

Do Not Use or Store Near Heat or Open Flame.

Rinse equipment and containers and dispose of wastes by burying. Destroy Empty Containers — Perforate, crush and bury in non-crop lands away from water supplies.

WARNING: Do not use this product on dairy animals, in dairy barns, or on feed or forage to be fed to dairy animals being finished for slaughter.

This product is poisonous to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

OBSERVE THE FOLLOWING PRECAUTIONS

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

DIRECTIONS

The following suggestions are general recommendations. For timing, number of applications and specific recommendations applicable to your area or crops, consult your State Agricultural Extension Service. Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use, as injury to plants may result.

RECOMMENDATIONS

Dosages are given in terms of pints of Toxaphene 6 Lb. Emulsion Concentrate per acre for use with sufficient water to provide thorough coverage.

ALFALFA: Armyworms, Cutworms, Grasshoppers, Leafhoppers, Lygusbugs, Thrips — Use 2 pints per acre. Apply in early spring or after cutting before new growth is 4 inches tall. Do not feed treated forage to dairy animals or animals being finished for slaughter.

CORN: Armyworms, Fall Armyworms. Use 2 pts. per acre when larvae are young. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on use of grain.

SMALL GRAINS (Barley, Oats, Wheat, Rye): Armyworms, Fall Armyworms. Use 2 pts. per acre when larvae are young. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restrictions on use of grains.

EPA Reg. No. 226-39

S 9-71

PESTICIDE

TOBACCO STATES CHEMICAL CO., INC.

ACCEPTED
June 5, 1972
UNDER THE
FUNGICIDE
FOR FEED
EPA REG. NO. 226-39