

Tobacco States


ACCEPTED
MAR - 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is registered under EPA Reg. No. 228-40

Lice & Tick Powder

CONTAINS 1% LINDANE

FOR CONTROL OF LICE, TICKS, FLEAS and MANGE attacking BEEF CATTLE, HORSES and PIGS



ACTIVE INGREDIENTS:	% by Wt
Gamma isomer of Benzene Hexachloride from Lindane	1.00%
INERT INGREDIENTS:	99.00%
TOTAL	100.00%

Keep Out Of Reach Of Children
CAUTION
See Other Cautions On Side Panel

PRECAUTIONARY STATEMENTS

CAUTION

Statement of Practical Treatment

- IF SWALLOWED** - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
- IF INHALED** - Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN** - Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES** - Flush eyes with plenty of water. Call a physician if irritation persists.

HAZARD TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. May be absorbed through skin. Avoid breathing of dust. Avoid contamination of feed or foodstuffs. Avoid dusting feeding troughs or utensils. May cause irritation of nose and throat. May be absorbed through skin. Applicators must wear the following protective clothing during the application process: a light weight protective suit or coveralls, water-resistant hat, unlined waterproof gloves, and unlined, light-weight boots. Mixers and loaders must also wear goggles or a face shield, waterproof gloves and a waterproof apron. Wear a disposable paper dust mask which covers at least one third of the face. This product should be applied in a well ventilated area. Avoid prolonged breathing of spray, dust, or vapor mist. Do not have any part of skin. Wash thoroughly after using and before eating or smoking.

STORAGE: Do not store in areas of extreme heat or cold. Protect from moisture or extreme humidity.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of empty pesticide containers, or residues in a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty can into application equipment. Then dispose of empty can in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this Product in a manner inconsistent with its labeling. (Do not apply within 30 days of slaughter)

- BEEF CATTLE** - Cattle Lice and Fleas: Dust the entire animal thoroughly. Repeat applications when necessary. Clean and dust stables and bedding. Keep untreated animals away from the treated cattle.
- SHEEP and GOATS** - Sheep-Ticks and Lice: Dust the entire animal thoroughly and repeat applications in 3 or 4 weeks if any live ticks or lice are found at that time. Do not use on dairy goats.
- PIGS** - Lice, Fleas, Ticks & Mange: Dust animals thoroughly and repeat applications in 14 days. Treatment of bedding, pens or houses is necessary.
- PONIES and MULES** - Lice: Dust coat of animal thoroughly. Do not rub in, but leave on top of coat.
- DOGS** - Lice, Fleas, and Ticks: Dust dogs with this material along the length of the back and work the powder in well. It is essential to clean and dust bedding on living quarters.