

DIRECTIONS — DILUTE WITH WATER

TOBACCO — Wireworms: Use 1 to 2 fluid ounces in 50 gallons of transplanting water, depending on the type of soil. Apply at the rate of 1 pint to the plant or about 200 to 250 gallons of this mixture to the acre. This establishes a zone of protection around the roots of the newly set plants allowing them to become well established. Note: Do not apply this product to crops in rotation with root crops and peanuts.

LIVESTOCK APPLICATIONS

NOTE: Be sure finished sprays of Lindane are thoroughly mixed at the time of application. This is important in order to avoid animal injury and to assure uniform effectiveness. Do not apply exaggerated dosages. Repeat applications only as often as necessary. Do not treat animals under 3 months of age. Do not treat sows within two weeks of farrowing nor at least three weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals. Do not spray animals within 30 days of slaughter. Do not dip animals within 60 days of slaughter. Do not use on dairy animals.

BEEF CATTLE, HOGS, TICKS, LICE, HORN FLY, STABLE FLY, MOSQUITOES: Use Dilution Table No. B, treating the entire animal and infested areas for insect control. For STABLE FLY, this treatment is only supplementary and should be accompanied by the treatment for FARM BUILDINGS AND BARNs housing these animals.

SHEEP AND GOATS: Treat in fall of year for sheep ticks or sarcoptic mange, use 2 quarts of Dilution Table No. B applying 2 gallons per adult animal at about 400 lbs. pressure. Do not spray animals within 30 days of slaughter. Do not use on dairy goats.

For a dip to control lice and sheep ticks use Table No. A. Keep the vat agitated at all times, completely submerge the animal's body including the head at least twice. Do not dip overheated animals. Always water animals before dipping so they won't drink the dipping solution. Do not dip animals within 60 days of slaughter.

FOR RESIDUAL FLY CONTROL ON AGRICULTURAL PREMISES SUCH AS Barns, Sheds, and Animal Shelters, fences (Excluding dairy barns or in poultry houses) — House Fly and Stable Fly: Use Dilution Table No. C Applying at the rate of one gallon of spray per 500 sq. ft. thoroughly wetting all surfaces to point of run-off. Protect all feed, feed troughs before spraying. Remove animals from barn. Repeat as required.

NOTE: Do not use treated manure on lands where root crops are to be planted.

DILUTION TABLE
Quantities of Water
Quantity of 20% Lindane To Use

	TABLE A	TABLE B	TABLE C
5 gal.	8 tsp.	10 tsp	9 fl. oz
25 gal.	6 fl. oz.	pt.	3 pts

2 Tablespoonfuls equal 1 fluid ounce.
3 Teaspoonfuls equal 1 tablespoon.

Do not use inside dairy barns, milk rooms or milk processing rooms. Do not use on dairy cattle, dairy goats, around feed or forage for animals or on animals being finished for slaughter. NEVER USE at rate stronger than recommended. Do not use on pets. Do not use in dwellings.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke).

This product is poisonous to fish and wildlife. Keep out of any body of water.

Apply this product only as specified on this label.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 226-68

S2-68

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY

ACCEPTED WITH COMMENTS

