

NET WEIGHT ONE POUND

Tobacco States

# 25% Lindane\* WETTABLE POWDER

### ACTIVE INGREDIENTS:

Gamma isomer of Benzene Hexachloride (from Lindane) 25%

INERT INGREDIENTS: 75%

**WARNING. MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. MAY PRODUCE IRRITATION OF SKIN AND EYES. Avoid breathing of vapor, dust or spray mist. Do not get in eyes or on skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Avoid spraying feeding troughs or utensils.**

**CAUTION:** Keep out of reach of children. See back panel for additional cautions.

25% LINDANE WETTABLE POWDER should be used only at dosages given below or follow the recommendations of the State Agricultural Experiment Station

**NOTE:** Avoid the use of this product on food crops when there is a possibility of interference with flavor or residue remaining at harvest. Avoid excessive use of this product where soil in which root crops are to be planted may become contaminated and cause off-flavor to such crops. Do not feed treated foliage to dairy animals or animals being finished for slaughter.

Do not contaminate milk, milking equipment, feed or drinking water. Do not use inside dairy bars, milk rooms, or milk processing rooms. Do not use on dairy cattle, dairy goats, around feed or forage for animals being finished for slaughter. Never use at a rate stronger than recommended. Do not use on pets. Do not use in dwellings.

USDA Reg. No. 226-74

H 5-6C

ACCEPTED

June 17, 1968

226-74

**FOR RESIDUAL CONTROL OF ROACHES, ANTS, FLIES AND MOSQUITOES ON AGRICULTURAL PREMISES SUCH AS BARN, SHEDS, ANIMAL SHELTERS (Excluding dairy barns or dairies houses).**  
**FOR FLIES AND MOSQUITOES:** (As a Spray) Use 1 lb. to 30 gallons of water and apply at the rate of about 1 gallon of spray mixture per 500 sq. ft. of surface. Apply as a coarse spray to walls, ceilings, beams, screens, window frames, partitions, and similar places where flies congregate. Protect all food, feed troughs before spraying. Remove all animals from barn. Repeat as required.

(As a Dust) Use 1 lb. to 1 gallon of water. Dust around windows, doors, ceiling beams, and other places where flies gather. Repeat as needed. See NOTE above.

**FOR ROACHES AND ANTS:** Use 1 pound in 10 gallons of water. Spray thoroughly all basements, cracks, floors and other places where insects hide, also spray any insects present.

**LIVESTOCK:** Do not treat sows within two weeks of farrowing nor for at least three weeks thereafter. Avoid treating animals in cold, stormy weather, as injury may result from chilling. Avoid treating overheated or sick animals. Do not use on dairy animals.

**BEEF CATTLE:** For LICE AND TICKS on BEEF cattle, use 1 pound in 100 gallons of water. Spray the entire animal or their infested area. For SCAB OR SARCOPTIC MITES on BEEF cattle, use 1 lb. per 100 gallons of water, and using a power sprayer producing about 400 pounds pressure, apply 2 gallons of spray to the animal. Good control requires complete coverage and for a complete cleanup, two applications a week or 10 days apart are required. Young calves, especially those under 3 months of age may be injured by treatment. It is sometimes advisable however, to treat young calves, apply only 1 quart of diluted spray per animal. Do not apply later than 30 days before slaughter, or dip animals within 60 days of slaughter. Do not dip calves under 3 months of age.

**HOGS:** For hog louse and sarcoptic mange mite on hogs, use 1 lb. in 100 gallons of water. Spray thoroughly making a single repeat application to entire animal or infested area a week or 10 days later if necessary. Do not spray within 30 days of slaughter, or dip within 60 days of slaughter. Do not treat animals under 3 months of age.

**SHEEP AND GOATS:** For sheep ticks, lice and scab mite (scabies), use as a spray at the rate of 1 lb. in 100 gallons of water. Thorough wetting of the animal is essential for good control. A second spraying may be necessary 2 weeks after the first application. As a dip, use 1 lb. per 100 gallons of water. A single dipping is often or usually sufficient for control. Treat in the fall. In some areas a spring spraying after shearing may be advisable. Do not spray within 30 days of slaughter. Do not dip within 30 days of slaughter. Do not use on dairy goats. Do not treat animals under 3 months of age.

**TICKS AND CHIGGERS ON VEGETATION:** Use 2 pounds per 100 gallons of water when ticks or chiggers are present. Thoroughly wetting vegetation. Repeat as necessary.

Use of all dosages listed below are based upon the amount of 25% Lindane required in 100 gallons of water.

**COLE CROPS (Cabbage, Broccoli, etc.):** For APHIDS, THRIPS, FLEA BEETLES, CABBAGE LOOPERS, IMPORTED CABBAGE WORM: Use 1 pound. Do not apply after heads begin to form.  
**CUCURBITS (Cucumber, Squash):** For CUCUMBER BEETLES AND THRIPS, spray with 1 lb. per 100 gallons of water. Repeat as required. For MELON AND PICKLE WORMS use 1 lb. Repeat application at 10 day to two week intervals. Treat both upper and lower leaf surfaces. Do not apply closer than 1 day of harvest.

**STRAWBERRIES:** For weevil and spottedwing, use 1 pound in 500 gallons of water when insects appear. Do not apply after fruit starts to form.

**NURSERY AND ORNAMENTAL PLANTS:** For aphids, spray with 1 lb.

**SEED TREATMENT:** For the temporary protection of germinating seeds and seedlings from attack by WIREWORM AND SEED MAGGOT: Dust the amounts listed below on each 100 pounds of seed at the time of planting. **CORN:** use 5 oz. per bushel or 1 lb. per 2 bushels. **BEANS:** 1 to 2 oz. per bushel. **MILK AND PEAS:** 5 oz. per bushel. Do not use seed as food or feed. Treated seeds exposed on soil surfaces may be hazardous to birds and other wildlife.

**Tobacco:** In "Setter Water" for wireworm. Mix 1 oz. of 25% Lindane Wettable Powder in each 10 gallons of water keeping the water agitated at rather frequent intervals, using about 1 qt. of the base of each plant, approximately 250 gallons of water per acre.

Rinse equipment and containers and dispose of wastes by burning in non-occupied lands away from water supplies. Containers should be disposed of by punching holes in them and burning with wastes or by burning (Keep out of smoke).

This product is poisonous to fish and wildlife. Keep out of any body of water.

Apply this product only as specified on this label. Tobacco States Chemical Co. Inc. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statements on this label.

PESTICIDE

TOBACCO STATES CHEMICAL CO., INC.  
LEXINGTON, KENTUCKY