

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn bee-keepers well in advance to locate hives at a safe distance until one week after application. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

**DIRECTIONS FOR USE**

READ ENTIRE LABEL USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

**DIRECTIONS—Do Not Dilute**

Vegetable Dust may be applied when first signs of insect infestation occur, preferably before damage from feeding occurs. Repeat as needed at 7 to 10 day intervals. For best results apply thoroughly to upper and lower leaf surfaces. Do not feed treated vines to dairy or meat animals.

Crops	Insects Controlled	Recommended Dose Per Acre	Time Between Last Application & Harvest
Beans	Mexican bean beetle	10 lbs.	1 Day before harvest
	leafhoppers, beet	20 lbs.	
	leaf-beetle, cucumber beetle, western bean cutworm, flea beetles and Japanese beetle.	20 to 25 lbs.	
	Corn earworm, armyworms, velvet bean caterpillar	40 lbs.	
Cucumbers Summer squash	Pickleworm and melonworm	10 lbs.	1 Day before harvest
	Cucumber beetles, squash bugs and flea beetles	20 lbs.	
Sweet Corn	Corn earworm, European corn borer, fall armyworm, and beetles and Japanese beetle	30 to 40 lbs.	1 Day before harvest for fresh market ears, 7 days before harvest on forage and husks fed to livestock
Tomatoes Eggplant Peppers	Tomato fruitworm, fall armyworm, flea beetles and leafhoppers	20 lbs.	1 Day before harvest
	Tomato hornworm, European corn borer and stink bugs	40 lbs.	
Potatoes	Colorado potato beetle	10 lbs.	1 Day before harvest
	Leafhoppers and flea beetle	20 lbs.	
	Colorado bugs	40 lbs.	

For insects attacking silks and ears, apply at 2 to 3 day intervals beginning when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation.

**CAUTION: HARMFUL IF INHALED OR SWALLOWED.**


Avoid prolonged or repeated breathing of dust. Do not take internally. SKIN CONTACT MAY BE HARMFUL. Avoid skin Contact. Wash Hands and Face Before Eating.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.


AVOID CONTAMINATION OF FOOD. KEEP OUT OF REACH OF CHILDREN.

USDA Reg. No. 226-147

S-11-68



Tobacco States Pesticides




NET CONTENTS—2 POUNDS



**VEGETABLE DUST**  
CONTAINS 5% SEVIN\*



For Control of Certain Insect Pests of Vegetable



**CAUTION:**  
Keep out of reach of children.

See other cautions on rear panel.

Active Ingredient: Carbaryl (1-naphthyl N-methylcarbamate) \_\_\_\_\_ 9% by wt.  
Inert Ingredients \_\_\_\_\_ 91% by wt.  
\*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.



Between Last  
tion & Harvest  
before harvest

before harvest

before harvest  
ish market  
days before  
on forage  
usks fed to  
ck  
tervals beginning  
ry. Three or more  
ifestation.

before harvest

before harvest


Internally.  
Wash Hands and  
ole cholinesterase

8/11, 1971  
UNDER THE FEDERAL INSECTICIDE  
ACT AND RODENT ACT  
ECONOMIC POISON REGISTERED  
EPA REG. NO. \_\_\_\_\_ SUBJECT  
REMARKS/COMMENTS.

226-147


# Tobacco States Pesticides

NET CONTENTS--2 POUNDS



## VEGETABLE DUST

CONTAINS 5% SEVIN\*



For Control of Certain Insect Pests of Vegetable



**CAUTION:**

**Keep out of reach of children.**

See other cautions on rear panel.

Active Ingredient:  
Carbaryl (1-naphthyl N-methylcarbamate) \_\_\_\_\_ 9% by wt.  
Inert Ingredients \_\_\_\_\_ 95% by wt.  
\*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

TOBACCO STATES CHEMICAL CO., INC.  
LEXINGTON KENTUCKY