



Net Contents

5 Pounds

# 7 1/2 % SEVIN<sup>®</sup> DUST



## For Control of Certain Insect Pests of Beans

**Active Ingredient:**

Carbaryl (1-naphthyl N-methylcarbamate) . . . 7 1/2% by wt.

Inert Ingredients . . . . . 92 1/2% by wt.

<sup>®</sup>Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn bee-keepers well in advance to locate hives at a safe distance until one week after application.

**CAUTION HARMFUL IF INHALED OR SWALLOWED**

Avoid prolonged or repeated breathing of dust. Do not take internally. **SKIN CONTACT MAY BE HARMFUL.** Avoid Skin Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Avoid prolonged or repeated breathing of dust. Do not take internally. **SKIN CONTACT WITH ANY INSECTICIDE MAY BE HARMFUL.** Avoid Unnecessary Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work.

Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.**

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

**CAUTION:** Keep out of the reach of children.

See other cautions on rear panel

USDA Reg. No. 226-170

**DIRECTIONS FOR USE**  
**READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.**

**DIRECTIONS—Do Not Dilute**

7 1/2% Sevin Dust may be applied when first signs of insect infestation occur, preferably before damage from feeding occurs. Repeat as needed at 7 to 10 day intervals. For best results apply thoroughly to upper and lower leaf surfaces. Do not feed treated vines to dairy or meat animals.

Crops	Insects Controlled	Recommended Dose Per Acre	Time Between Last Application & Harvest
Beans	Mexican bean beetle	7 lbs.	1 Day before harvest
	Leafhoppers, bean leaf beetle, cucumber beetles, western bean cutworm, flea beetles and Japanese beetle.	13 1/3 lbs.	
	Corn earworm, armyworms, velvet bean caterpillar	13 1/3 to 17 lbs.	
	Lygus and stink bugs	27 lbs.	

Tobacco States Chemical Co., Inc. **MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED,** concerning this product or its use, which extend beyond the statements on this label.

Do not reuse empty containers. Destroy them by burying with waste or burning. Stay away from smoke or fumes.

5-11-67

