

Tobacco States Chemical Co.

MANUFACTURERS AND DISTRIBUTORS
INSECTICIDES, FUNGICIDES, AND WEED KILLERS

P. O. BOX 479

Lexington, Kentucky 40501

PHONE: 606-233-1446

PRODUCT DATA SHEETS

Name of product: Tomahawk Brand Super X-Termite
EPA Reg. No. 226-177

Uses: None other than stated on label

MIXING DIRECTIONS OR TERMITE PROOFING & CONTROL OF SUBTERRANEAN TERMITES

Mix the contents of this container with 48 parts of water, or at the rate of 5 1/3 tablespoons of this concentrate per gallon of water. Stir thoroughly to get good mixture of concentrate and water. Apply this mixture as follows:

DIRECTIONS FOR TERMITE PROOFING & CONTROL OF SUBTERRANEAN TERMITES

NOTICE: Soil Poisoning and Void Poisoning are very necessary to Termite Proof Buildings. Termites must have contact with the moisture and woody tissue. The necessary moisture is usually obtained from the soil. Wood treated in place does not take up enough chemical to make it completely resistant to termites. Hence it is important to get both soil moisture and kill termites that have contacted the soil. Make annual inspections for possible reinfestation.

BUILDINGS HAVING CRAWL SPACES

- (A) Dig a trench adjacent to and around all piers and pipes and along both sides of the foundation walls. Dig the trench to, but not below the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons of mixed Super X-Termite per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons of mixture per 10 linear feet.
- (B) Treat voids in hollow-block masonry foundations at the rate of 1 gallon of mixture per 5 linear feet of wall. Apply so that the chemical will reach the footing. If this is done by drilling or probing rod holes AVOID GOING INTO PLUMBING OR ELECTRICAL CONDUITS.

BUILDINGS HAVING BASEMENTS

- (A) Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundations, dig a trench to, BUT NOT BELOW, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons of mixed Super X-Termite per 10 linear feet FOR EACH FOOT OF DEPTH. A trench 3-feet deep would require 12 gallons of mixture per 10 linear feet.
- (B) It may also be necessary to treat critical areas only under the basement flooring such as around sewer pipes, conduits and piers and along the inside of the foundation walls and interior walls. One method consists of drilling holes about a foot apart through the concrete floor and adjacent to the areas requiring treatment. The mixed Super X-Termite should be injected into the soil beneath the floor. AVOID DRILLING INTO PLUMBING OR ELECTRICAL CONDUITS. The mixture should be applied at the rate of at least 4 gallons per 10 linear feet of wall.

(Directions continued on other panel)

21076

NET CONTENTS: 1 GAL.

**TOMAHAWK
BRAND
SUPER X-TERMITE**

"DO-IT-YOURSELF" CONCENTRATE
for TERMITE CONTROL

ACTIVE INGREDIENTS:	
Technical Chlordane*	45.00%
Gamma Isomer Benzene	
Hexachloride from Lindane	1.00%
Heavy Aromatic Naphtha	48.00%
INERT INGREDIENTS:	
	6.00%

*Equivalent to 27.00% Octachloro-4,7-methanotetrahydroindane and 18.00% of related compounds.

CAUTION: Keep out of reach of children.
See other Cautions on back panel

BELKNAP, INC. LOUISVILLE, KENTUCKY 40202

- (C) Treat voids in hollow-block foundations at the rate of 1 gallon per 5 linear feet of wall so that the mixture will reach the footing. Do this by drilling or probing. TAKE CARE TO AVOID DRILLING INTO PLUMBING OR ELECTRICAL CONDUITS.

SLAB-ON-GROUND CONSTRUCTION

- (A) Infestations in this type of construction are difficult to control. One method consists of drilling holes about a foot apart through the concrete slab, adjacent to all cracks and expansion joints, and injecting mixed Super X-Termite into the soil beneath the slab. AVOID DRILLING INTO PLUMBING OR ELECTRICAL CONDUITS. Another method is to drill through the foundation walls from the outside and force the mixture just beneath the slab along the inside of the foundation and along all the cracks and expansion joints. The mixture should be applied at the rate of at least 4 gallons per 10 linear feet of foundation or expansion joint.

- (B) Dig a trench 1 foot in depth, but not below the top of the footing, along the outside of the foundation walls. Apply the mixture at the rate of 4 gallons per 10 linear feet of trench. Mixed Super X-Termite should be mixed with the soil as it is being replaced in the trench.

- (C) Treat voids in hollow block foundations at the rate of one gallon of mixture per 5 linear feet of wall so that the chemical will reach the footing. Do this by drilling or probing. AVOID DRILLING INTO PLUMBING OR ELECTRICAL CONDUITS.

CAUTION: This concentrate and the mixtures from this solution may cause toxic symptoms if brought into contact with the skin. Avoid inhalation and skin contact. After using, and in case of spillage on the skin, wash immediately with soap and water. Harmful if swallowed. Keep out of reach of children. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CAUTION: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Do not reuse empty container. Destroy when empty by perforating or crushing, bury or discard with wastes or by burning (keep out of smoke).

Belknap, Inc., makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

EPA Reg. No. 226-177

EPA Est. 226-KY-1

ACCEPTED
SEP 7, 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC PEST CONTROL

DIRECTIONS Continued (28-4)

BROWN DOG TICKS—Use 2% spray solution (4 oz. Riverdale Chlordane 4 in 3 quarts water). Thoroughly

treat infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets and localized areas of floor and floor coverings. Use fresh bedding for pets. Repeat applications as necessary. Do not use on pets.

CARPET BEETLES—Use 2% spray solution (4 oz. Riverdale Chlordane 4 in 3 quarts water). Thoroughly treat infested area of carpets, localized areas of floor and baseboards and cracks under carpets where insects may be found. Repeat as necessary. Note: Use as spot treatment to areas listed.

SPIDERS AND CENTIPEDES—Use 2% spray solution (4 oz. Riverdale Chlordane 4 in 3 quarts water). Thoroughly treat infested baseboards, corners, pipes, storage localities and other infested areas in basements. Repeat as needed.

HOUSE FLIES, MOSQUITOES AND WASPS—Use 2% spray solution (4 oz. Riverdale Chlordane 4 in 3 quarts water). For indoor application spray thoroughly and repeatedly doors, around windows and areas frequented by these insects. For outdoor application to aid in preventing entry, spray screens, window frames, door sills and selected areas where these insects tend to alight or gather. Repeat applications as necessary. For wasps, spray the nest with a double strength solution (8 oz. in 3 quarts water) after dark when all the insects have returned to the nest.

Riverdale CHLORDANE 4

Contains 4 lbs. Technical Chlordane Per Gallon

ACTIVE INGREDIENTS:

Technical Chlordane*

Petroleum Distillates

INERT INGREDIENTS

By Weight

45.0%

51.0%

4.0%

TOTAL 100.0%

*Equivalent to 27% Octachloro-4,7-Methanotetrahydroindane and 18% related compounds

CAUTION: KEEP OUT OF REACH OF CHILDREN See additional Warnings on Side Panel

DIRECTIONS FOR USE

SHADE TREES, ORNAMENTALS, PERENNIALS AND ANNUALS—To control bag worms, box elder bugs, elm leaf beetles, leafminers, lace bugs, tree crickets, thrips, Japanese Beetle Adults, June Beetle Adults, use 2 quarts in 100 gallons water. Apply spray when insects are first observed covering plant thoroughly until runoff appears. For smaller quantities mix 2 teaspoonsful in 1 gallon water.

LAWN AND ORNAMENTAL TURF—Ants—Mix 4 teaspoons in 1 gallon water or 1 quart in 50 gallons water. Apply to colony, thoroughly soaking area. Ticks, Chiggers—Mix 3 pints Riverdale Chlordane 4 in 100 gallons water per acre. Apply to lawn and shrubbery. For smaller quantities mix 1/12 pint per gallon water to cover 1,000 sq. ft.

Green June beetle larvae, Japanese beetle larvae, White Fringed Beetle larvae, White Grubs, Wireworms—Mix 2 1/2 - 5 quarts in 100 gallons water per acre. Wash into the top few inches of soil. If dosage is doubled to 5 to 10 quarts it will give a minimum of 2 to 3 years control.

Mosquito Control—For control of adult mosquitoes mix 6 gallons in 100 gallons water and apply 4 gallons of dilute liquid per acre. Apply to grass and shrubbery. Applications may be made to walls and ceilings of porches and to out-buildings. For smaller amounts mix 6 teaspoons to 1 gallon water. Do not use in dairy barns and poultry houses.

FIELD CROPS: CORN—For control of corn rootworm apply 3 quarts in necessary water to cover 1 acre. Make application prior to or at time of planting, covering with soil immediately. For row application use 2 quarts. To control wireworms, white grubs, seed corn beetles, seed corn maggots, Japanese Beetle grubs and cutworms increase dosage to 4 quarts and apply as broadcast treatment.

OATS, RYE—To control armyworm mix 1 quart in sufficient water to cover one acre. Do not spray oats or rye after heads start to form. Do not graze treated crop.

HOUSEHOLD APPLICATION—Mix 4 oz. of Riverdale Chlordane 4 in 3 quarts water. This forms an effective spray solution (2%) to control: Ants, crickets, roaches, silverfish and waterbugs. Thoroughly spray cracks and crevices, insect trails, hiding places and areas around door sills, window frames, openings around water pipes, heat ducts, electrical outlets, baseboards and other locations where insects might enter. Spray on insects if possible. Repeat applications as necessary.

CAUTION:

Harmful if swallowed. This solution may be absorbed through skin. Therefore, avoid contact with skin, eyes and clothing. Do not breathe spray mists. In case of spillage on skin wash immediately. Avoid contamination of foodstuffs and feed. Keep out of reach of children. Do not use on household pets or humans. Do not spray near or toward an open flame. Do not smoke while spraying. Keep children and pets off treated lawns until spray solution has been washed into the soil by watering or rain and grass is thoroughly dry. This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Do not use in commercial food preparation areas or in edible products areas of food processing plants. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

E.P.A. Reg. No. 228-41

NET CONTENTS 1 qt.

RIVERDALE CHEMICAL COMPANY

1000 W. 10th St., Kansas City, Mo. 64101