

NET CONTENTS—5 POUNDS



# CHLORDANE 20 GRANULAR

Active Ingredient:	
*Technical Chlordane .....	20%
Inert Ingredients: .....	80%
	100%

\*Equivalent to 12% Octachloro-4, 7-methanotetrahydroindane and 8% Related Compounds

### CAUTION:

- KEEP OUT OF REACH OF CHILDREN
- HARMFUL IF SWALLOWED.
- AVOID BREATHING DUST.
- AVOID CONTACT WITH SKIN.
- WASH WITH SOAP AND WATER AFTER EXPOSURE.
- AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

See Additional Precautions on Back



### CAUTION:

This product is toxic to man, birds, and other animals, and to other wildlife in treated areas may be present. Do not apply where birds are likely to occur. Do not contaminate water, food, or feed, or equipment, or disposal of wastes. Apply this product only as specified on the label.

### ANTIDOTE:

EXTERNAL: In case of spillage on skin, wash promptly with soap and water.

INTERNAL: Emetic of One (1) Teaspoon of mustard in a Tumbler of water. Call a Physician.

### CONTAINER DISPOSAL:

Do not reuse empty containers. Reuse equipment if it is torn and dispose of wastes by carrying out to dump and away from water supplies. Containers should be destroyed by punching holes in them and burning with waste by burning. Keep out of smoke.


### DIRECTIONS FOR USE

CORN SOIL INSECTS: For control of corn rootworms apply a broadcast treatment at the rate of 12 1/2 lbs. per acre after the field is plowed. For control of Japanese Beetles apply 1 1/2 lbs. per acre August 1, 1976.

INSECT CONTROL ON NURSERY SITES For Use Under U.S.D.A. Japanese Beetle Quarantine Programs Only. For the control of Japanese Beetles on Nurseries where ornamental plants are or will be at the time of application of Chlordane 20 Granular apply 3 lbs. per acre of granular to cover surface uniformly using fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat ornamentals with wet. Apply before rain or sprinkling.

EPA Reg. No. 106-178

EPA Est. No. KY-1



TOBACCO STATES CHEMICAL CO.  
LEXINGTON KENTUCKY

NET CONTENTS—5 POUNDS



# CHLORDANE 20

# GRANULAR

Active Ingredient:	
*Technical Chlordane .....	20%
Inert Ingredients: .....	80%
	<hr/>
	100%

\*Equivalent to 12% Octachloro-4, 7-methanotetrahydroindane and 8% Related Compounds

### CAUTION:

- KEEP OUT OF REACH OF CHILDREN
- HARMFUL IF SWALLOWED.
- AVOID BREATHING DUST.
- AVOID CONTACT WITH SKIN.
- WASH WITH SOAP AND WATER AFTER EXPOSURE.
- AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

See Additional Precautions on Back



### CAUTION:

This product is toxic to fish, birds, and other wildlife in treated areas only. Do not apply where runoff of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### ANTIDOTE:

**EXTERNAL:** In case of spillage on skin, wash promptly with soap and water.

**INTERNAL:** Emetic if One (1) tablespoon of mustard oil or Tumbler of water. Call a Physician.

### CONTAINER DISPOSAL:

Do not reuse empty containers. Place equipment in a safe place and dispose of wastes by burying in a deep and away from water supply. Containers should be disposed of by punching holes in them and burning in the yard or by burning. Keep out of reach.


### DIRECTIONS FOR USE

**CORN SOIL INFESTED:** For control of larvae, this granule is a first class treatment. Apply 100 lbs per acre in the field after the field has been plowed and before the corn is planted. Effective August 1st.

**INSECT CONTROL IN HOME GARDENS:** For Use Under U.S.D.A. Japanese Beetle Quarantine Programs Only. For the control of Japanese Beetles, apply this granule where ornamental plants are grown. Use at the rate of 10 lbs of Chlordane 20 Granular per acre. For control of Japanese Beetles to cover surface of lawn, use 10 lbs per acre. Treatment should be applied in the spring and fall. For control of Japanese Beetles on lawns, use 10 lbs per acre. Apply before rain or irrigation.

EPA Reg. No. 10124

EPA EST. NO. 1



TORRICO STATES CHEMICAL

**CAUTION:**

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**ANTIDOTE:**

**EXTERNAL:** In case of spillage on skin, wash promptly with soap and water.

**INTERNAL:** Emetic of One (1) Tablespoon of mustard in a Tumbler of water. Call a Physician.

**CONTAINER DISPOSAL:**

Do not reuse empty containers. Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Keep out of smoke.

**DIRECTIONS FOR USE**

**CORN SOIL INSECTS:** For control of Cutworms apply as first dust treatment at the rate of 12 1/2 to 20 lbs. per acre after the field is plowed. Use as promptly. Use suspended effective August 1, 1976.

**INSECT CONTROL ON NURSERY SOILS. For Use Under U.S.D.A. Japanese Beetle Quarantine Programs Only.** For the control of Japanese Beetle Grubs, in Nurseries where ornamental plants are grown, use at the rate of 25 lbs. of Chlorfume 20 Granular per acre (2/3 lb. to each 1,000 sq. ft.) to cover surface uniformly using a fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling.

EPA Reg. No. 226-178

EPA Est. 26KY1

**ACCEPTED**

JUN 2 1976

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 226-178

TORRICO STATES CHEMICAL CO

NEW YORK KENTUCKY

TASCO

TASCO  
BRAND