

ACCEPTED

MAR 30 1944

LIBRARY OF THE
BUREAU OF AERONAUTICS
WASHINGTON, D. C.

CROP	TO CONTROL	DIRECTIONS	
		DOSAGE LBS./100 GALS	APPLICATION
Ornamentals (Greenhouse and Out of Doors)	Aphids	1	On ornamentals...
	Cyclamen Mite		
CAUTION: This product may injure birch trees and certain varieties of magnolia trees such as Bonnafor Deluxe.			
Strawberries	Cyclamen Mite	1	Spray plants thoroughly. Apply . . . gallons of spray per acre and thoroughly wet stems, foliage, and crowns of the plants. For multiple applications, do not apply at intervals less than 35 days during the fruiting period. Do not apply within 4 days of harvest.
	Meadow Spittlebug Strawberry Aphid	1	Apply as required. Do not apply at intervals of less than 35 days or more than twice during any 35 day period when fruit is present. Do not apply within 4 days of picking berries.
Tobacco	Flea Beetle	1	Seed Beds: Apply about 6 gallons per 100 square yards. Field: Use 1 to 2 pounds per acre and apply in sufficient water for uniform coverage. Do not apply within 5 days of harvest.
	Hornworm Budworm Green Peach Aphid		
Broccoli Cabbage	Aphid	1 1/2 to 2	Do not apply within 7 days of harvest.
	Cabbage Worm Flea Beetle Harlequin Bug Stink Bug		
Eggplants Tomatoes (Field and Greenhouse)	Blister Beetle	1	Do not apply within 1 day of harvest.
	Colorado Potato Beetle Flea Beetle Green Peach Aphid Green Stink Bug Whitefly Hornworm	1 to 2	
Peppers	Flea Beetle	1	Do not apply within 1 day of harvest.
	Colorado Potato Beetle Leafhopper Aphid		
Cucumbers	Aphid	1 to 2	For vine borer control, apply weekly to flower buds, stems and vines beginning when moths first appear. May be applied to harvest.
Mellons Cucumbers	Cucumber Beetle		
Pumpkins	Squash Vine Borer		
Squash	Striped Cabbage		
	Flea Beetle		

CAUTION: Workers entering treated areas on the day of applications should wear protective clothing.