



Net Contents 3 Pounds

95% FERBAM

WETTABLE POWDER
AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

Ferbam (Ferric Dimethyldithiocarbamate) 95%

INERT INGREDIENTS 5%


CAUTION: Keep Out of Reach of Children.

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID BREATHING DUST OR SPRAY MIST. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION. "Do not contaminate or store near feed or foodstuffs."

"This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label."

USDA Reg. No. 226-189

TOBACCO STATES CHEMICAL CO., INC.
Lexington, Kentucky



DIRECTIONS FOR USE AS A SPRAY

All Recommendations Based on 100 Gallons of Spray

- APPLES:** No applications later than 7 days before harvest.
 - Scab — ¾ to 1½ pounds at pre-bloom, calyx and early-cover sprays; ½ pound later cover sprays. In combination with sulphur use ¾ pound plus 3 pounds wettable sulphur. Not recommended for Golden Delicious and other varieties susceptible to russeting.
 - Rust — Use scab recommendations through first cover and include a special bloom spray of ¾ pound.
 - Other Diseases — Use scab recommendations as long as any one or more of the following diseases may appear — Apple Blotch, Bitter Rot, Black Rot, Blossom End Rot, Brooks Spot, Fly Speck, Frog-eye Leaf Spot and Sooty Blotch.
- ALMONDS:** Shot-Hole — 1¼ to 1½ pounds at popcorn and petal-fall.
- APRICOTS:** No applications later than 21 days before harvest.
 - Brown Rot and Shot-Hole — 1½ pounds at red-bud, early bloom and full-bloom.
 - Green or Jacket Rot — 1½ pounds at full-bloom and petal-fall.
- BLUEBERRIES:** No applications later than 40 days before harvest.
 - Mummy Berry — 2¼ pounds at loose bud scales and 7 days later.
 - Botrytis Blight — 2¼ pounds immediately before bloom, full-bloom and 2 to 3 weeks later.
- CABBAGE:** Downy Mildew in Plant Beds — 1½ pounds 2 to 3 times a week beginning when plants emerge.
- CELERY:** Early Blight (Cercospora), Late Blight (Septoria) and Damping-Off — ¾ to 1½ pounds once a week with first application in the plant bed and continuing in the field. Under severe conditions spray every 3 to 5 days. Remove residues by stripping, trimming and washing.
- CHERRIES:** Leaf Spot — Apply 1½ pounds 3 times at 14 day intervals starting at petal-fall. Spray foliage again immediately after harvest.
 - Brown Rot — 1½ to 1½ pounds at pink bud stage, full-bloom and petal-fall and at 7 to 10 day intervals starting 2 to 3 weeks before harvest. A spreader may be added to late sprays to avoid blotching fruit.
- CITRUS (Florida) Scab:** Apply 1¼ lbs. at pre-bloom and ¾ petal-fall and late summer or early fall if heavy flush of growth occurs.
- CRANBERRIES:** No applications later than 28 days after mid-bloom.
 - Fruit Rots — 2¼ pounds 3 times at 14 day intervals with first spray at mid-bloom.
 - Fairy Ring Disease — 7½ lbs. in the fall immediately after harvest. Apply one gallon per square foot to the diseased area and up to 3 feet outside of the advancing ring of dead plants.
- GRAPES:** No applications later than 7 days before harvest.
 - Black Rot — 1½ pounds just before bloom, immediately after bloom and 14 days after bloom. Additional applications may be made on recommendation of State Experiment Station or State Extension Service authorities at delayed dormant, cluster break, berry touch, and 10 to 14 days after berry touch stages.
- LETTUCE:** Damp-Off in Plant Beds — ¾ pound as a drench (4 to 5 gallons per 100 sq. ft.) when plants emerge and at 14 day intervals.
- PEACHES:** No application within 21 days of harvest.
 - Peach Leaf Curl and Shot-Hole — 1½ to 2¼ pounds at dormant and popcorn.
 - Brown Rot Blossom — 1½ to 1½ pounds at pink bud and 25 percent to 75 percent of full bloom.
 - Fusicoccum Canker — 1¼ pounds at 7 to 10 day intervals before bloom and after harvest.
- PEARS:** No applications later than 7 days before harvest.
 - Scab — 1¼ pounds in pink, calyx, first and second-cover sprays.
 - Fabaea and Sooty Blotch — ¾ pound in summer sprays.
- PINE SEEDLINGS:** Fusiform Rust in Plant Beds — 1½ pounds with suitable spreading agent. Apply at 3 to 7 day intervals from plant emergence as needed.
- PLUMS AND PRUNES:** No applications later than 7 days before harvest.
 - Brown Rot — 1¼ to 1½ pounds at popcorn, full-bloom and 2 weeks later. Apply again 5 weeks before harvest and repeat at 7 day intervals during wet weather.
 - Leaf Spot — ¾ to 1¼ pounds in fall after harvest and before leaf drop. Repeat in spring as buds swell.
- RASPBERRIES, BLACKBERRIES, BOYSENBERRIES:** No applications later than 40 days before harvest.
 - Anthracoze, Rust and Spur Blight — 2¼ pounds delayed dormant, 1¼ to 2 pounds when new canes are 10 to 12 inches high and before bloom.
- TOBACCO:** Blue Mold — 2½ to 3 pounds at 3 to 4 day intervals beginning when leaves are size of a dime and continue until transplanting time. Use 3 gallons of suspension per 100 square yards of bed when plants are small, increasing to 6 gallons as plants enlarge.
- TOMATOES:** Gray Mold and Sclerotinia (Florida) — 1½ to 3 pounds at 3 to 5 day intervals beginning at transplanting time. Apply in combination with other fungicides used for control of other diseases.