

Tobacco States Chemical Co.

MANUFACTURERS AND DISTRIBUTORS
INSECTICIDES, FUNGICIDES, AND WEED KILLERS

P. O. BOX

PHONE (804) 233-1446


NET CONTENTS 10 OZ.



MANEB 80

WITH ZINC ADDED

80% WETTABLE POWDER AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENTS:	
Maneb (Manganese Ethylenebisdithiocarbamate)	80%
(Total Manganese as metallic	16.5%)
INERT INGREDIENTS:	20%
	100%

CAUTION: Keep out of reach of children.
See back panel for additional precautions.

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID BREATHING DUST OR SPRAY MIST. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION. Keep away from fire and sparks.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

EPA Reg. No. 226-190

M 8-71

DIRECTIONS FOR USE AS A SPRAY

TOMATOES:

Early Blight, Late Blight, Septoria Leaf Spot, Gray Leaf Spot (Stemphylium) and Anthracnose — 4 tablespoons per 3 gallons of water applied when fruit of first clusters is well formed and repeated at 5 day intervals. (To avoid damage, do not use on tender young plants in greenhouse.) Under severe weather conditions favoring disease, apply more frequently beginning when plants are first set in fields. Do not apply within 5 days of harvest.

POTATOES:

Early and Late Blights — 4 tablespoons per 3 gallons of water applied when plants are 2 to 6 inches high and repeating at 5 to 10 day intervals as long as blight threatens. Under severe conditions use the shorter interval.

ONIONS:

Downy Mildew, Purple Blotch (Blight) and Blast (Botrytis Leaf Blight) — 4 tablespoons per 3 gallons of water applied when first spots appear, repeating weekly as required.

LETTUCE:

Downy Mildew — 4 tablespoons per 3 gallons of water. Apply when disease appears, repeating at 10 day intervals under average conditions and 3 to 5 day intervals when disease conditions are severe. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce by washing or other effective means.

Watermelons — Downy Mildew, Anthracnose, Cercospora Leaf Spot and Gummy Stem Blight — 4 tablespoons per 3 gallons of water. Apply when plants begin to run or when disease threatens, repeating at 5 day intervals as required. Shorten interval between application to 3 to 5 days under severe conditions. Do not apply within 5 days of harvest.

CABBAGE:

Alternaria Leaf Spot and Downy Mildew — Apply 4 tablespoons per 3 gallons of water.

Plant Beds and Direct Seeded Fields — Apply 7 to 10 days after planting or earlier if disease is present. Repeat at 3 day intervals until plants are set in field or until field-seeded plants are thinned. Fields — Apply when the disease threatens; repeat weekly as needed. Shorten interval to 3 to 5 days under severe conditions. Do not apply later than 7 days before harvest.

CARROTS:

Leaf Spot Diseases — Use 4 tablespoons per 3 gallons of water. Apply when plants are 5 weeks old or when disease threatens, repeating every 7 to 10 days.

TOBACCO STATES CHEMICAL CO INC