

JASCO**MANEB 80****WITH ZINC ADDED****80% WETTABLE POWDER
AGRICULTURAL FUNGICIDE**

ACTIVE INGREDIENTS:	% By Wt.
Maneb. (Manganese Ethylenebisdithiocarbamate) ..	80.00%
(Total Manganese as metallic -----)	16.50%
INERT INGREDIENTS: -----	20.00%
TOTAL -----	100.00%

KEEP OUT OF REACH OF CHILDREN**CAUTION****STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED — Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED — Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN — Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES — Rinse for at least 15 minutes with water and call a physician immediately.

**SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA Reg. No. 226-190

EPA Reg. No. 226-190

TOBACCO STATES CHEMICAL CO., INC.

LEXINGTON, KENTUCKY

PRECAUTIONARY STATEMENTS

CAUTION**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. During the mixing and loading of this product, please wear the following protective clothing: long pants, long sleeve shirt, gloves, hat and boots. Do not enter treated area without protective clothing until dusts have settled. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. In case of contact, flush with plenty of water. For eyes, get medical attention. Keep away from fire and sparks.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

STORAGE & DISPOSAL

STORAGE: Do not store in areas of extreme heat or cold. Protect from moisture or extreme humidity.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag or can into application equipment. Then dispose of empty bag or can in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this Product in a manner inconsistent with its labeling.

TOMATOES: Early Blight, Late Blight, Septoria Leaf Spot, Gray Leaf Spot (Stemphylium) and Anthracnose — 4 tablespoons per 3 gallons of water applied when fruit of first clusters is well formed and repeated at 5 day intervals. (To avoid damage, do not use on tender young plants in greenhouse.) Under severe weather conditions favoring disease, apply more frequently beginning when plants are first set in fields. Do not apply within 5 days of harvest.

POTATOES: Early and Late Blights — 4 tablespoons per 3 gallons of water applied when plants are 2 to 6 inches high and repeating at 5 to 10 day intervals as long as blight threatens. Under severe conditions use the shorter interval.

ONIONS: Downy Mildew, Purple Blotch (blight) and Blot (Botrytis Leaf Blight) — 4 tablespoons per 3 gallons of water applied when first spots appear, repeating weekly as required.

LETTUCE: Downy Mildew — 4 tablespoons per 3 gallons of water. Apply when disease appears, repeating at 10 day intervals under average conditions and 3 to 5 day intervals when disease conditions are severe. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce by washing or other effective means.

WATERMELONS — Downy Mildew, Anthracnose, Cercospora Leaf Spot and Gummy Stem Blight — 4 tablespoons per 3 gallons of water. Apply when plants begin to run or when disease threatens, repeating at 5 day intervals as required. Shorten interval between application to 3 to 5 days under severe conditions. Do not apply within 5 days of harvest.

CABBAGE: Alternaria Leaf Spot and Downy Mildew — Apply 4 tablespoons per 3 gallons of water.

Plant Beds and Direct Seeded Fields — Apply 7 to 10 days after planting or earlier if disease is present. Repeat at 3 day intervals until plants are set in field or until field-seeded plants are thinned. **Fields** — Apply when the disease threatens; repeat weekly as needed. Shorten interval to 3 to 5 days under severe conditions. Do not apply later than 7 days before harvest.

CARROTS: Leaf Spot Diseases — Use 4 tablespoons per 3 gallons of water. Apply when plants are 6 weeks old or when disease threatens, repeating every 7 to 10 days.

WARRANTY STATEMENT

Tobacco States **MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED,** concerning this product or its use, which extend beyond the statements on this label.

226-190
PM 2/1
Pg. 2 of 1