

TOBACCO STATES BRAND

ROSE DUST


ACTIVE INGREDIENTS
 Zineb, Zinc ethylene bisdithiocarbamate
 Methoxychlor, Technical
 Gummi, solvent of benzene hexachloride, 100%
 Wettable
 Inert Ingredients
 Methoxychlor, Technical
 Wettable
 Gummi, solvent of benzene hexachloride, 100%
 Wettable

FOR ROSES & ORNAMENTALS

ALSO USED AS A SPRAY

CAUTION

Keep out of Reach of Children. See other Cautions on back panel.



ROSE DUST

Begin application before diseases are apparent and before insects become numerous. Protect plants through the growing season by dusting or spraying at 7 to 10 day intervals and after heavy rains. Cover thoroughly the (upper and lower) leaf surfaces and stems with a light film of dust or spray. Applications should be in the late afternoon while temperature is low and wind is still. Note: Do not apply dust while foliage is wet.

AS A SPRAY: Mix 8 tablespoons to 1 gallon of water.


PARTIAL LIST OF INSECTS CONTROLLED:

Aphids or plant lice, Rose Chafer, Rose Leaf Beetle, Rose Leafhopper, Thrips, Rose Slugs, Red Spider Mites, Clover Mite, Flea Beetles, Grasshoppers, Caterpillar, (tent and puss), Cucumber Beetle and certain other leaf eating insects.

PARTIAL LIST OF DISEASES CONTROLLED:

Roses - Black Spot, Powdery Mildew, Rust.
 Other Flowers - such as (Aster, Carnation, Chrysanthemum, Begonia, Belladonna, Dahlia, Calendula, Delphinium, Hollyhock, Hydrangea, Snapdragon, African Violet, Lilac, Zinnia, Azaleas, Camellias, Larkspur) - Powdery Mildew, Rust, Leaf Spot, Anthracnose.
 Gladioli - Botrytis, Curvularia, Stemphylium.

CAUTION: Harmful if swallowed. May be absorbed through skin. Avoid prolonged breathing of dust or spray mist, avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.



USDA Reg. No. 226-196



TOBACCO STATES CHEMICAL CO. INC. LEXINGTON, KENTUCKY

ACTIVE INGREDIENTS:

Zineb (Zinc ethylene bisdithiocarbamate)	6.0%
*Methoxychlor, Technical	5.0%
Gamma isomer of benzene hexachloride from lindane	1.0%
2-(1-Methylheptyl)-4,6-dinitrophenyl crotonate9%
Other nitrophenols and derivatives, chiefly 1-methylhepty-4,6-dinitrophenol1%

INERT INGREDIENTS 87.0%

*5% Methoxychlor, Technical is equivalent to 2,2-Bis
(p-methoxyphenyl)-1,1,1-trichloroethane 4.4% and
related compounds 0.6%.

**R ROSES &
AMENTALS**

Begin application before diseases are apparent and before insects become numerous. Protect plants through the growing season by dusting or spraying at 7 to 10 day intervals and after heavy rains. Cover thoroughly the