


UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 22,222 SUBJECT  
TO ATTACHED COMMENTS.

NET CONTENTS-4 POUNDS



Tobacco States  
Pesticides

# 5 % SEVIN VEGETABLE DUST



For Control of Certain Insect Pests of Vegetable

Active Ingredients:  
Carbaryl (1-naphthyl N-methylcarbamate) ..... 5% by wt.  
Inert Ingredients ..... 95% by wt.  
\*Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

## CAUTION:

Keep out of reach of children.  
See back panel for additional cautions.

TOBACCO STATES CHEMICAL CO.  
LEXINGTON  
KENTUCKY

**DIRECTIONS FOR USE**  
**READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL**  
**DIRECTIONS AND CAUTIONS.**

**DIRECTIONS—Do Not Dilute**

5% Sevin Dust may be applied when first signs of insect infestation occur preferably before damage from feeding occurs. Repeat as needed at 7 to 10 day intervals. For best results apply thoroughly to upper and lower leaf surfaces. Do not feed treated vines to dairy or meat animals.

| Crops                                                                                                                                                                                                                                                                                                                                                              | Insects Controlled                                                                                                                                                                                                                                                                                                    | Recommended<br>Dose<br>Per Acre                                                                            | Time Between Last<br>Application & Harvest                                                                                                                                                                                                                                                              |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Beans                                                                                                                                                                                                                                                                                                                                                              | Mexican bean beetle<br>Leafhoppers, bean<br>leaf beetle, cucumber<br>beetles, western bean<br>cutworm, flea beetles<br>and Japanese beetle.<br>Corn earworm,<br>armyworms, velvet<br>bean caterpillar<br>Lygus and<br>stink bugs                                                                                      | 10 lbs.<br>20 lbs.<br><br><br>20 to 25 lbs.<br><br>40 lbs.                                                 | 1 Day before Harvest                                                                                                                                                                                                                                                                                    |
| Cucumbers<br>Summer<br>squash                                                                                                                                                                                                                                                                                                                                      | Pickleworm and<br>melonworm<br>Cucumber beetles,<br>squash bugs and flea<br>beetles                                                                                                                                                                                                                                   | 10 lbs.<br>20 lbs.                                                                                         | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Sweet Corn                                                                                                                                                                                                                                                                                                                                                         | Corn earworm,<br>European corn<br>borer, fall army-<br>worm, sap beetles<br>and Japanese beetle                                                                                                                                                                                                                       | 30 to 40 lbs.                                                                                              | 1 Day before harvest<br>for fresh market<br>ears, 7 days before<br>harvest on forage<br>and husks fed to<br>livestock                                                                                                                                                                                   |
| For insects attacking silks and ears, apply at 2 to 3 day intervals beginning when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation. "Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population." |                                                                                                                                                                                                                                                                                                                       |                                                                                                            |                                                                                                                                                                                                                                                                                                         |
| Tomatoes<br>Eggplant<br>Peppers                                                                                                                                                                                                                                                                                                                                    | Tomato fruitworm,<br>fall armyworm,<br>flea beetles and leaf-<br>hoppers<br>Tomato hornworm,<br>European corn<br>borer and stink bugs                                                                                                                                                                                 | 20 lbs.<br>40 lbs.                                                                                         | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Potatoes                                                                                                                                                                                                                                                                                                                                                           | Colorado potato<br>beetle<br>Leafhoppers and flea<br>beetle<br>Lygus bugs                                                                                                                                                                                                                                             | 10 lbs.<br>20 lbs.<br>40 lbs.                                                                              | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Poultry<br>(Chickens,<br>Turkeys,<br>Ducks, Geese,<br>Game Birds,<br>Pigeons)                                                                                                                                                                                                                                                                                      | ON BIRDS: Northern<br>fowl mite, chicken<br>mite, lice, fleas<br>IN PREMISES:<br>Chicken mite,<br>fleas, bedbugs,<br>fowl tick<br>ON FLOOR LITTER:<br>Lesser mealworms,<br>northern fowl mite,<br>chicken mite, lice,<br>fleas, bedbugs<br>DUST BATH BOXES:<br>Northern fowl mite,<br>chicken mite, lice<br>and fleas | 1 lb. per<br>100 birds<br><br><br>1 lb. per<br>40 sq. ft.<br><br>2 1/2 lb.<br>per box for<br>each 50 birds | 7 days before slaughter.<br>Avoid contamination<br>of nests, eggs, feed,<br>and water troughs.<br>Tolerance 5 ppm.<br>on meat and fat, 0<br>in eggs. When dusting<br>poultry, do not repeat<br>application within<br>4 weeks. Do not house<br>birds in treated houses<br>within 7 days of<br>slaughter. |
| Dogs and<br>cats                                                                                                                                                                                                                                                                                                                                                   | Brown dog ticks,<br>fleas                                                                                                                                                                                                                                                                                             | Rub in skin<br>and apply in                                                                                | Don't treat kittens<br>until 4 weeks old.                                                                                                                                                                                                                                                               |

|                                                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                       |                                                                                                        |                                                                                                                                                                                                                                                                                                         |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                                                                                                                                                                                                                                                                                    | bean caterpillar<br>Lygus and<br>stink bugs                                                                                                                                                                                                                                                                           | 40 lbs.                                                                                                |                                                                                                                                                                                                                                                                                                         |
| Cucumbers<br>Summer<br>squash                                                                                                                                                                                                                                                                                                                                      | Pickleworm and<br>melonworm<br>Cucumber beetles,<br>squash bugs and flea<br>beetles                                                                                                                                                                                                                                   | 10 lbs.<br>20 lbs.                                                                                     | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Sweet Corn                                                                                                                                                                                                                                                                                                                                                         | Corn earworm,<br>European corn<br>borer, fall army-<br>worm, sap beetles<br>and Japanese beetle                                                                                                                                                                                                                       | 30 to 40 lbs.                                                                                          | 1 Day before harvest<br>for fresh market<br>ears, 7 days before<br>harvest on forage<br>and husks fed to<br>livestock                                                                                                                                                                                   |
| For insects attacking silks and ears, apply at 2 to 3 day intervals beginning when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation. "Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population." |                                                                                                                                                                                                                                                                                                                       |                                                                                                        |                                                                                                                                                                                                                                                                                                         |
| Tomatoes<br>Eggplant<br>Peppers                                                                                                                                                                                                                                                                                                                                    | Tomato fruitworm,<br>fall armyworm,<br>flea beetles and leaf-<br>hoppers<br>Tomato hornworm,<br>European corn<br>borer and stink bugs                                                                                                                                                                                 | 20 lbs.<br>40 lbs.                                                                                     | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Potatoes                                                                                                                                                                                                                                                                                                                                                           | Colorado potato<br>beetle<br>Leafhoppers and flea<br>beetle<br>Lygus bugs                                                                                                                                                                                                                                             | 10 lbs.<br>20 lbs.<br>40 lbs.                                                                          | 1 Day before harvest                                                                                                                                                                                                                                                                                    |
| Poultry<br>(Chickens,<br>Turkeys,<br>Ducks, Geese,<br>Game Birds,<br>Pigeons)                                                                                                                                                                                                                                                                                      | ON BIRDS: Northern<br>fowl mite, chicken<br>mite, lice, fleas<br>IN PREMISES:<br>Chicken mite,<br>fleas, bedbugs,<br>fowl tick<br>ON FLOOR LITTER:<br>Lesses mealworms,<br>northern fowl mite,<br>chicken mite, lice,<br>fleas, bedbugs<br>DUST BATH BOXES:<br>Northern fowl mite,<br>chicken mite, lice<br>and fleas | 1 lb. per<br>100 birds<br><br>1 lb. per<br>40 sq. ft.<br><br>2 1/2 lb.<br>per box for<br>each 50 birds | 7 days before slaughter.<br>Avoid contamination<br>of nests, eggs, feed,<br>and water troughs.<br>Tolerance 5 ppm.<br>on meat and fat, 0<br>in eggs. When dusting<br>poultry, do not repeat<br>application within<br>4 weeks. Do not house<br>birds in treated houses<br>within 7 days of<br>slaughter. |
| Dogs and<br>Cats                                                                                                                                                                                                                                                                                                                                                   | Brown dog ticks,<br>fleas                                                                                                                                                                                                                                                                                             | Rub in skin<br>and apply in<br>sleeping<br>quarters weekly                                             | Don't treat kittens<br>until 4 weeks old.                                                                                                                                                                                                                                                               |

CAUTION: HARMFUL IF INHALED OR SWALLOWED.

Avoid breathing of dust. Do not take internally.

SKIN CONTACT MAY BE HARMFUL. Avoid skin contact. Wash Hands and Face Before Eating.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD.

KEEP OUT OF REACH OF CHILDREN.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn bee-keepers well in advance to locate hives at a safe distance until one week after application. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

USDA Reg. No. 226-207