

NET CONTENTS ONE GALLON

Tobacco States

No. 2 Formula

Six

FOR CONTROL OF

1. Aphid
 2. Flea Beetle
 3. Hornworm
 4. Cabbage Looper
 5. Budworm
 6. Grasshopper
- ON TOBACCO**

TOBACCO

SPRAY

ACTIVE INGREDIENTS

Malathion (0,0-dimethyl phosphorodithioate diethyl mercaptosuccinate)	22.320%
Endosulfan (Hexachlorocyclopentadiene-2,4,3-benzodioxathiepin oxide)	14.880%
Heptachlor Epoxide (heptachlor)	56.600%
INERT INGREDIENTS	6.200%

*Incl. in U.S. Pat. No. 2,799,685 Reg. Off. of Canadian Her. Inst. 11/1/55

WARNING: Keep out of reach of children.

See other Warnings on back panel.

USDA REG NO. 225,211

§ 173

WARNING: Harmful if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water for 15 minutes. Medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Bureau of Mines for endosulfan protection. Do not allow to drift to areas occupied by unprotected humans or beneficial animals. Do not contaminate or store near feed or foodstuffs.

NOTE TO PHYSICIANS: Malathion upon repeated, prolonged use may cause cholinesterase inhibition. Atropine is antidote to PAM also antidotal and may be administered in conjunction with atropine. Inrodon produces central nervous system excitation and responds to the barbituric acid derivatives.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur or where it may cause damage to crops, water by cleaning of equipment, or disposal of wastes.

Containers should be disposed of by punching holes in them and burying with waste.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES, OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

DIRECTIONS FOR USE — DILUTE WITH WATER — (Follow Directions)

SIX-XX TOBACCO SPRAY No. 2 Formula is an emulsifiable insecticide formulation designed for use as a preventative spray for the control of six major insect pest on tobacco. This formulation can be used in any type sprayer. If mixture is allowed to stand, re-agitate before use.

TOBACCO - field, green peach aphid, hornworms, budworms - use 1 to 2 quarts per acre. For leaf beetles and small cabbage looper - use 2 quarts per acre. Grasshoppers - Use 3 quarts. Dosage rates given should be used in sufficient water to give adequate coverage depending on the type of nozzle and equipment being used and in the size of the crop. Cover the entire plant very lightly with insecticide. Do not soak plants, as injury may occur. Do not apply in the heat of the day as some burning of plants could result. When using hand sprayers hold the nozzle at least 12 inches above the plant. Do not direct the spray on the bud as injury may result. Check fields every 3 to 5 days, and repeat as needed depending on the severity of the infestation and local conditions. Do not apply within 6 days of harvest.

For plants up to knee high use approximately 1½ gallons of water per acre. Plants taller use 300 gallons. For best control of named insects, use a preventative program of spraying early in the season when the plants are small. Larvicide or insecticides on leaves must be sprayed at the best results. For weeds, frequently for reinfestations.

For any and all applications, the required amount of spray will be 2 to 5 gallons of water per acre. Insects entering the field heads within 24 hours after application should wear protective clothing.

DO NOT STORE NEAR HEAT OR OPEN FLAME DO NOT STORE BELOW 20 °F

THIS STATEMENT OF WARRANTY DOES NOT EXTEND TO ANY OTHER STATES OR COUNTRIES. THE MAKER MAKES NO WARRANTY OR MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE EXPRESS OR IMPLIED CONCERNING THE PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE STATEMENTS ON THIS LABEL.

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON KENTUCKY

ACCEPTED

File # 1971

UNDER THE FEDERAL JOBS
TRAINING ACT, TO DE
FOR ECONOMIC REHAB
ED UNDER 2022 C 211
U. S. ATTORNEY GENERAL