




Net contents 25 lbs.

BUDWORM DUST

INSECTICIDES CONTROLS FLEA BEETLES IN PLANT BEDS



ACTIVE INGREDIENT:

‡Endosulfan (Hexachlorohexahydromethano-2,1,3-benzodioxathiepin oxide) 3.0%

INERT INGREDIENTS: 97.0%
100.0%

‡Thiodan U.S. Pat. No. 2,799,685

USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking, wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated fields within 24 hours should wear protective clothing.

USDA Reg. No. 226-211

DIRECTIONS FOR USE

Apply the listed amounts per acre when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe days interval between last application and harvest indicated by number in () following crop.

TOBACCO (5): Cabbage Looper, Green Peach Aphid, Tobacco Budworm, Tobacco Flea Beetle, Tobacco Hornworm, PLANT BED: Use 2 lbs. per 100 sq. yds. FIELD: Use 20 to 35 lbs. per acre.

For Budworms:

Examine buds in several areas of field for feeding signs and small green or tan caterpillars, lightly treat top of plants. Check plants and re-treat after 10 days if needed. Do not apply within 75 days of harvest.

Allow 3 days to kill budworms. DO NOT REPEAT WITHIN 30 DAYS.

TOBACCO STATES CHEMICAL CO., INC.
LEITCHVILLE, KENTUCKY