


TOMATO

Contains
Zineb



Contains
Methoxychlor

MULTI-PURPOSE DUST



ACTIVE INGREDIENTS:
 Zineb (Zinc ethylene bisdithiocarbamate) 5.0%
 (Metallic Zinc equivalent 1.23%)
 Methoxychlor Technical 5.0%

INERT INGREDIENTS 90.0%
 5.0% Methoxychlor, technical is equivalent to
 4.4% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and .6% related compounds.

CAUTION: Keep out of reach of children.

Avoid breathing dusts. Avoid contamination of feed and food-stuffs. Do not use on food crops later than 7 days of harvest; except on broccoli, do not use within 14 days of harvest. Remove all residue from food crops by washing, brushing or other effective means of harvest. This product is poisonous to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE — Do Not Mix With Water

TOMATO MULTI-PURPOSE DUST may be applied to moist or dry foliage any time when dusting conditions are favorable. Begin applications soon after plants are above ground or when disease conditions threaten or insects first appear. Apply at 5 to 10 day intervals or more often if disease conditions become severe. Use 20 to 50 lbs. of dust per acre (1/4 to 1 pound per 1,000 square feet) depending on the crop, size of plants, and severity of disease or insect conditions.

CROPS

INSECTS & DISEASES

TOMATOES PEPPERS

Flea Beetle, Leafhopper, Fruitworm, Fall Armyworm, Suck Fly, Blister Beetle, Early and Late Blight, Anthracnose, Stemphylium and Septoria Leaf Spot, Cladosporium Gray Mold: Begin spraying when disease is reported in area and repeat application at 7 to 10 day intervals or oftener if blight conditions are severe. Do not apply within 7 days of harvest.

BEANS

Mexican Bean Beetle, Corn Earworm, Leafhopper, Downy Mildew, Rust, Anthracnose: Begin when first blossoms appear. Repeat application at 4 to 7 day intervals. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI & CAULIFLOWER

Cross-striped Cabbage Worm, Diamond Back Moth, Imported Cabbage Worm, Flea Beetle, Downy Mildew, Alternaria Leaf Blight: Do not apply after harvest. Do not apply within 7 days of harvest; except broccoli, do not apply within 14 days of harvest.

CANTALOUPE, CUCUMBERS, WATERMELONS

Cucumber Beetle (various), Leafhopper, Flea Beetle, Squash Vine Borer, Downy Mildew, Alternaria, (Macrosporium) Blight, Anthracnose, Gummy Stem Blight: Direct spray especially at under-side of leaves, begin applications when plants appear above the ground. Do not apply within 7 days of harvest.

CHRYSANTHEMUMS, CARNATIONS, HOLLY-HOCKS, SNAPDRAGONS

Rust Leaf Spot, Anthracnose, Leafhopper.

NOTE: Under severe weather conditions where control of diseases specified is necessary, continue at weekly intervals with Dithane spray or dust alone. Remove excess residues from edible parts by washing, brushing or other effective means at harvest.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

USDA Reg. No. 226-216

H 6 71

TOBACCO STATES CHEMICAL CO. INC.