

Tobacco States Chemical Co., Inc.

FORMULATOR AND DISTRIBUTOR
AGRICHEMICALS AND SPRAY EQUIPMENT

PHONE: (606) 233-1445

P. 4

NET CONTENTS ONE POUND (453.6 gms)



or Dust"

ACCEPTED
FEB 1 1963
Under the Federal
Fungicide, and
as amended, is
registered and
EPA Reg. No. 226-216

ACTIVE INGREDIENTS:

Zineb (Zinc ethylene bisdithiocarbamate) 5.0%
(Metallic Zinc equivalent 1.23%)
Methoxychlor Technical 5.0%

INERT INGREDIENTS:

..... 90.0%
5.0% Methoxychlor, technical is equivalent to
4.4% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloro-
ethane and .6% related compounds.

CAUTION: Keep Out of Reach of Children

Avoid breathing dusts. Avoid contamination of feed and food.
stuffs.

*Additional
Clothing
Caution
HERE.*

*Text will be
in boldface
on new label.*

This product is poisonous to fish. Keep out of lakes, streams,
and ponds. Apply this product only as specified on this label.
This product is toxic to bees and should not be applied when
bees are actively visiting the area.

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY

NET CONTENTS ONE POUND (453.6 gms)



ACTIVE INGREDIENTS:
 Zineb (Zinc ethylene bisdithiocarbamate) 5.0%
 (Metallic Zinc equivalent 1.23%)
 Methoxychlor technical 5.0%

INERT INGREDIENTS: 90.0%
 5.0% Methoxychlor, technical is equivalent to
 4.4% 2,2-bis (p-methoxyphenyl)-1, 1, 1-trichloro-
 ethane and .6% related compounds.

CAUTION: Keep Out of Reach of Children

Avoid breathing dusts. Avoid contamination of feed and food stuffs.

This product is poisonous to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Add protective clothing cautions here.

Text will be in boldface on new label.

DIRECTIONS FOR USE — Do Not Mix With Water

TOMATO MULTI-PURPOSE DUST may be applied to moist or dry foliage any time when dusting conditions are favorable. Begin applications soon after plants are above ground or when disease conditions threaten or insects first appear. Apply at 5 to 10 day intervals or more often if disease conditions become severe. Use 20 to 50 lbs. of dust per acre (½ to 1 pound per 1,000 square feet) depending on the crop, size of plants, and severity of disease or insect conditions.

CROPS	INSECTS & DISEASES
TOMATOES PEPPERS	Flea Beetle, Leafhopper, Fruitworm, Fall Armyworm, Bister Beetle, Early and Late Blight, Anthracnose, Stemphylium and Septoria Leaf Spot, Cladosporium Gray Mold. Begin dusting when disease is reported in area and repeat application at 7 to 10 day intervals or oftener if blight conditions are severe.
BEANS	Mexican Bean Beetle, Corn Earworm, Leafhopper, Downy Mildew, Rust, Anthracnose. Begin when first blossoms appear. Repeat application at 4 to 7 day intervals.
CABBAGE, BROCCOLI & CAULIFLOWER	Cross-striped Cabbage Worm, Diamond Back Moth, Imported Cabbage Worm, Flea Beetle, Downy Mildew, Alternaria Leaf Blight.
CANTALOUPE, CUCUMBERS, WATERMELONS	Cucumber Beetle (various), Leafhopper, Flea Beetle, Squash Vine Borer, Downy Mildew, Alternaria (Macrosporium) Blight, Anthracnose, Gummy Stem Blight. Direct dust especially at under-side of leaves begin applications when plants appear above the ground.
CHRYSANTHEMUMS, CARNATIONS, HOLLY-HOCKS, SNAPDRAGONS	Rust Leaf Spot, Anthracnose, Leafhopper.

NOTE: Under severe weather conditions where control of diseases specified is necessary, continue at weekly intervals with Dithane spray or dust alone. Remove excess residues from edible parts by washing, brushing or other effective means at harvest.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EPA Reg. No. 226-216
EPA Est. 226 - KY - 1

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY

MA22581

Tobacco States Chemical Co., Inc.

FORMULATOR AND DISTRIBUTOR
AGRICHEMICALS AND SPRAY EQUIPMENT

P. O. BOX 12530

Lexington, Kentucky 40583-2530

PHONE: (606) 233-1446

PRODUCT DATA SHEET

NAME OF PRODUCT: Tobacco States Brand "Tomato Copper Dust"
E.P.A. Reg. No. 226-32

CLAIMS: NONE OTHER THAN STATED ON LABEL.

1983
226-32

ACCEPTED
FEB 11 1983
Under the Federal Insecticide,
Fungicide and Rodenticide Act
for the pesticide
226-32

During the mixing and loading of this product, please wear the following protective clothing; long pants, long sleeve shirt, gloves, hat and boots.

ACCEPTED
FEB 11 1983
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
the pesticide
226-211

During the mixing and loading of this product, please wear the following protective clothing; long pants, long sleeve shirt, gloves, hat and boots.



1983 FEB 11 10 41 AM

NET CONTENTS ONE POUND (453.6 gms)



ACTIVE INGREDIENTS:

Zineb (Zinc ethylene bisdithiocarbamate) 50%
 (Metalic Zinc content 17.5%)
 Methoxychlor Technical 50%

INERT INGREDIENTS:

90.0%
 50% Methoxychlor (technical) and 44% Zinc ethylene bisdithiocarbamate and 6% related compounds

CAUTION: Keep Out of Reach of Children

Avoid breathing dusts. Avoid contamination of feed and food stuffs.

This product is poisonous to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE — Do Not Mix With Water

TOMATO MULTI-PURPOSE DUST may be applied to moist or dry foliage any time when dusting conditions are favorable. Begin applications soon after plants are above ground or when disease conditions threaten or insects first appear. Apply at 5 to 10 day intervals or more often if disease conditions become severe. Use 20 to 50 lbs. of dust per acre (1/2 to 1 pound per 100 square feet) depending on the crop, size of plants, and severity of disease or insect conditions.

CROPS	INSECTS & DISEASES
TOMATOES PEPPERS	Flea Beetle, Leafhopper, Fruitworm, Fall Armyworm, Blister Beetle, Early and Late Blight, Anthracnose, Stemphylium and Septoria Leaf Spot, Cladosporium Gray Mold. Begin dusting when disease is reported in area and repeat application at 7 to 10 day intervals or oftener if blight conditions are severe.
BEANS	Mexican Bean Beetle, Corn Earworm, Leafhopper, Downy Mildew, Rust, Anthracnose. Begin when first blossoms appear. Repeat application at 4 to 7 day intervals.
CABBAGE, BROCCOLI & CAULIFLOWER	Cross-striped Cabbage Worm, Diamond Back Moth, Imported Cabbage Worm, Flea Beetle, Downy Mildew, Alternaria Leaf Blight.
CANTALOUPE, CUCUMBERS, WATERMELONS	Cucumber Beetle (various), Leafhopper, Flea Beetle, Squash Vine Borer, Downy Mildew, Alternaria (Watermelon) Blight, Anthracnose, Gummy Stem Blight. Dust dust especially at underside of leaves. Begin applications when plants appear above the ground.
CYRUSPELLETS, CAPSICUMS, HOT PEPPERS, SNAPPEAGONS	Rust Leaf Spot, Anthracnose, Leafhopper.

ACCEPT
 FEB 11
 Under the Feds
 Fungicide, and
 as amended, to
 registered and
 EPA Reg No.

NOTE: Under severe weather conditions where control of diseases specified is difficult, combine at weekly intervals with D-thane spray or dust alone. Remove insects from edible parts by washing, brushing or other effective means at harvest.
 Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

EPA Reg No 226-216
 EPA No. 226 - KY - 1

TOBACCO STATES CHEMICAL CO., INC.
 LEXINGTON, KENTUCKY

Tobacco States Chemical Co., Inc.

FORMULATOR AND DISTRIBUTOR
AGRICHEMICALS AND SPRAY EQUIPMENT

P. O. BOX 12530

Lexington, Kentucky 40583-2530

PHONE: (606) 233-1446

PRODUCT DATA SHEET

NAME OF PRODUCT: Tobacco States Brand "Tomato Copper Dust"
E.P.A. Reg. No. 226-32

CLAIMS: NONE OTHER THAN STATED ON LABEL.

PTED
1 1983
226-32

