



DISINFECTANT

sanitizing animal housing and
to farrowing; sanitizing poul-
e bath for barns and poultry
itizing food-processing equip-

| | |
|----------------|----------------|
| ga-hydroxypoly | |
| x | 18.05% |
| iodine) | |
| | 16.00% |
| | 65.95% |
| Total | 100.00% |

following U. S. Patents:

| | |
|--|-----------|
| | 2,931,777 |
| | 2,977,315 |
| | 3,028,299 |

OUT OF REACH OF CHILDREN

ions on Back Panel.

No. 226-236

226-236 Comments

ACCEPTED

DIS-IN-FECT-IT DISINFECTANT Has A Built-in Indicator Of Germicidal Activity.
When the color disappears, a fresh solution should be prepared.

DIRECTIONS FOR USE

Gross filth must be removed prior to applications of the following recommended disinfecting solutions.

- FOR CLEANING AND DISINFECTING BUILDINGS AND EQUIPMENT ON THE FARM:**
Use 1/2 ounce per gallon water. Scrub surfaces to be cleaned and disinfected with a good brush or with a power sprayer. Start at the highest point of the building or equipment being cleaned, and work down. Let solution drain dry without rinsing.
- FOR CLEANING SOWS:**
Before placing sows in farrowing pens or crates, scrub down flanks and udder with a luke-warm solution of 1 ounce of disinfectant per gallon of water. Use a soft-bristled brush. Let sow drain dry and do not rinse until just before farrowing.
- SANITIZING POULTRY DRINKING WATER:**
Add 1 ounce of disinfectant to every 10 gallons of drinking water. The regular use in the drinking water also helps keep the waterers free of slime and mineral deposits.
- FOR SANITIZING BUILDINGS AND EQUIPMENT ON THE FARM:**
To sanitize surfaces of buildings, equipment or utensils which have previously been thoroughly cleaned, use 1 ounce of disinfectant in 5 gallons of water. Spray or swab solution on surfaces to be sanitized. Allow to drain dry without rinsing.
- SHOE BATH:**
To help prevent tracking disease organisms into poultry houses, farrowing houses and hog barns, place a shoe bath with 3 ounces of disinfectant per gallon of water inside the doorway. Scrape off shoes outside doorway and stand in shoe bath for 30 seconds prior to entering. Change shoe bath daily.
- DEODORANT:**
Dis-in-fect-it Disinfectant destroys many odors as it sanitizes and disinfects. For general deodorant applications (garbage pails,) a solution of 1 ounce of disinfectant per gallon of water.
- FOOD PLANT SANITIZATION:**
Use 1 ounce of Dis-in-fect-it Disinfectant in each 5 gallons of water as the final sanitizing rinse on previously cleaned food-processing equipment and utensils. Drain, but do not rinse. When used as so directed, Dis-in-fect-it Disinfectant does not require a final potable water rinse, as prescribed in Federal Food Additive Regulation 121.2547.

DANGER: Avoid contamination of food. Avoid contact with skin. Do not get in eyes. Causes eye damage or skin irritation. In case of contact, flush immediately with plenty of water. For eyes, get medical attention.

May be harmful if swallowed. If swallowed, do not induce vomiting. Drink milk, egg white, or plenty of water. Immediately get medical attention.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

TOBACCO STATES CHEMICAL CO., INC.

ACCEPTED WITH COMMENTS